


P.1
Donald R. Lynam, Ph.D., CIH, PE
July 6, 1990
Page 3

suggested, this should not stop us from introducing Mn to gasoline because the balance for and against doing so is clearly in favor of approving Ethyl's application.

Best regards.

Sincerely yours,

A handwritten signature in dark ink, appearing to read "H. Wisniewski", written over the typed name.

Henry M. Wisniewski, M.D., Ph.D.
Director

CURRICULUM VITAE

HENRYK M. WISNIEWSKI, M.D., Ph.D

DIRECTOR

Address: Home: 141 Nixon Avenue
Staten Island, NY 10304

Business: N.Y.S. Institute for Basic Research in Developmental
Disabilities

Birthdate: February 27, 1931, Luskowko, Poland
United States Citizen

Personal Status: Married, 2 Children

Education: M.D., Medical School, Gdansk (Danzig), Poland, 1955;
Ph.D., Medical School, Warsaw, Poland, Experimental
Pathology, 1960; Docent, Neuropathology, Medical School,
Warsaw, Poland, 1965.

Honors: Weil Award, American Association of Neuropathologists,
1969
Moore Award, American Association of Neuropathologists,
1972
Career Scientist, Health Research Council of the City
of New York
Association for the Help of Retarded Children, New York
City (AHRC) Chapter award, 1984
President, American Assoc. of Neuropathologists, 1984
Welfare League, Letchworth Village Chapter, AHRC, award
1985
Benevolent Society for Retarded Children, Staten Island
Chapter, 1986
Fellow, American Association for the Advancement of
Science, 1989

Experience: Director, NYS Institute for Basic Research in
Developmental Disabilities, 1976--present;
Professor of Pathology (Neuropathology), SUNY Health
Science Center at Brooklyn, 1976--present;
Director, MRC Demyelinating Diseases Unit, Newcastle
upon Tyne, England, 1974-1976;
Research Assoc., Assistant, Associate and full
Professor of Pathology (Neuropathology),
Albert Einstein College of Medicine, 1966-1975;
Visiting Scientist, Laboratory of Neuropathology
NINCDS, NIH (Dr. Klatzo), 1962-63;
Visiting Neuropathologist, Division of Neuropathology,
University of Toronto, Canada (Dr. J. Olszewski)
1961-1962;
Assistant-Associate Professor, Head of Laboratory of
Experimental Neuropathology, Associate Director
of Institute of Neuropathology, Polish Academy

of Science, Warsaw, Poland, 1958-1966.

Publications:

Author or co-author of 470 publications.

Societies:

American Association for the Advancement of Science
 American Association of Neuropathologists
 British Immunological Association
 British Society of Neuropathology
 Canadian Association of Neuropathologists
 American Association for Retarded Citizens
 Association for Research in Nervous and Mental Disease
 American Association on Mental Deficiency
 International Society for Developmental Neurosciences
 Society for Experimental Neuropathology
 Sigma Xi

Major Research
 Interests:

Developmental disabilities and dementia; Neuronal
 fibrous protein pathology; Demyelinating diseases.

Other Activities:

Past Member Neurology B Study Section, NIH. Ad hoc
 reviewer
 Past Member National Institute of Child Health and
 Human Development (NICHD), NIH, Mental
 Retardation Research Committee, Ad hoc reviewer
 Member of the Board of Directors of the Research
 Foundation for Mental Hygiene, Inc.
 Member of the Editorial Boards of:
 Acta Neuropathologica (Berl.)
 Neurotoxicology
 Journal of Neuropathology and Experimental
 Neurology
 Developmental Neuroscience
 Intl. Journal of Geriatric Psychiatry
 Alzheimer Disease & Associated Disorders
 International Journal
 Brain Dysfunction
 Dementia
 Member of the Journal of Neural Transmission -
 Dementia Section
 Member of the Editorial Advisory Board of Neurobiology
 of Aging
 Member of the Editorial Board of Brain Pathology

PUBLICATIONS

HENRYK M. WISNIEWSKI

1. Wisniewski, H., and Dobrzynska, Z. A case of the so called occult sclerosing carcinoma of the thyroid with metastases. Pol. Tyg. Lek. 26: 1009-1010, 1958.
2. Radzikowski, G., Ledochowski, A., Ledochowski, A., Wisniewski, H., and Nazarewicz Schwan, S. Research on anticarcinomatous substances. Patol. Pol. 9: 331-343, 1958.
3. Bittel-Dobrzynska, N., and Wisniewski, H. A case of hyperphosphatemia with kidney anomaly. Ped. Pol. 10: 1228-1234, 1958.
4. Osetowska, E., Gail, H., Lukasiewicz, D., Karcher, D., and Wisniewski, H.M. Leudocytrophie infantile precoce (type Krabbe). Rev. Neurol. 102: 63-477, 1960.
5. Saper, J., and Wisniewski, H. Brain stem tumor. Neurol. Neurochir. I Psych. Pol. 10: 713-716, 1960.
6. Wisniewski, H. The pathogenesis of some cases of cerebral hemorrhage. (A morphologic study of the margins of hemorrhagic foci and areas of the brain distant from the hemorrhage). Acta. Med. Pol. 2: 379-390, 1961.
7. Wisniewski, H. Research on experimental filling of the ventricular system of dogs. Acta Neuropath. 1: 238-244, 1961.
8. Wisniewski, W., Kruszewski, S., and Wisniewski, H. Radio-opaque media in pancreatography. Acta. Med. Pol. 229-235, 1961.
9. Nazarewicz, T., Ledochowski, Z., Konepa, J., Stenzel, J., Pikel, L., Falkowski, L., and Wisniewski, H. The study on the antineoplastic activity of poria oblique. Nowotwory 11: 401-411, 1961.
10. Wisniewski, H., and Gwozdziwicz, J. Three cases of a developmental anomaly of the nervous system with deformation of the Arnold-Chiari type. Pat. Pol. 12: 67-81, 1961.
11. Majewska, Z., Lehmanowa, J., Piolowski, J., Jankowski, T., and Wisniewski, H. Cerebral tumors in newborn and young children. Neurol. Neurochir. I Psychiat. Pol. 12: 7-14, 1962.
12. Wisniewski, H., Wrzolkowa, T., and Zyromska, M. Two cases of cerebellar medulloblastomas in adult patients. Neurol. Neurochir. I Psych. Pol. 12: 501-508, 1962.
13. Wisniewski, H., and Olszewski, J. Vascular permeability in the area postrema and hypothalamus. Neurol. 13: 885-894, 1963.
14. Wisniewski, H. Studies on atherosclerosis of the cerebral blood vessels. Neuropat. Pol. 2: 2-87, 1963.
15. Wisniewski, H. Hemorrhage and necrosis in the pituitary gland in cases with increased intracranial pressure, recent myocardial infarction and other pathological conditions. Neuropat. Pol. 3: 299-316, 1965.
16. Wisniewski, H., Majdecki, T., and Wisniewska, K. Topography of brain lesions after intracerebral, intraventricular and subarachnoid injection of copper salts. Neuropat. Pol. 3: 391-396, 1965.
17. Klatzo, I., Wisniewski, H., and Smith, D. Observations on penetration of serum proteins into the central nervous system. Prog. in Brain Res. 15: 73-88, 1965.

18. Klatzo, I., Wisniewski, H., and Streicher, E. Experimental production of neurofibrillary degeneration. J. Neuropath. Exp. Neurol. 24: 187-199, 1965.
19. Wisniewski, H. Studies on the permeability of the blood-brain, blood-cerebrospinal fluid and cerebrospinal fluid-brain barriers proteins under physiological conditions and of the blood-brain barrier for albumins in cerebral edema. Neuropat. Pol. 3: 1-33, 1965.
20. Streicher, E., Wisniewski, H., and Klatzo, I. Resistance of immature brain to experimental cerebral edema. Neurol. 15: 833-836, 1965.
21. Klatzo, I., Wisniewski, H., Steinwall, O., and Streicher, E. Dynamics of cold injury edema. Workshop on Brain Edema, Vienna, Sept. 11-13, 1965.
22. Osetowska, E., and Wisniewski, H. Ataxic familiale chez lapin de laboratoire diff. de la maladie de Sawin-Anders. Proc. of V Internat. Congr. Neuropath., Zurich, 1965.
23. Wisniewski, H., Pobudkowski, A., and Dembowski, J. Epileptic properties of phenol and cresol. Zwierzeta Lab. 3: 87-92, 1965.
24. Wisniewski, H., Smialek, M., Zalewska, T., Szydlowska, H., and Rap, Z. Copper contents in brains with blood-brain barrier damaged by freezing, hemorrhage and encephalomalacia. Neuropat. Pol. 4: 33-51, 1966.
25. Wisniewski, H., Karczewski, W., Wisniewska, K. Neurofibrillary degeneration of nerve cells after intracerebral injection of aluminum paste. Acta Neuropath. 6: 211-219, 1966.
26. Osetowska, E., Wisniewski, H., Wroblewska-Mularczyk, Z. Topography of tissue lesions and localization of the virus in tick-borne encephalitis. Pol Med. Journal 6: 134-150, 1966.
27. Wisniewski, H., Smialek, M., Szydlowska, H., and Zaleska, T. Quantitative topography of copper in Wilson's disease and in porto-systemic encephalopathy. Neuropat. Pol. 5: 91-103, 1967.
28. Wisniewski, H., Narkiewicz, O., and Wisniewska, K. Topography and dynamic of neurofibrillar degeneration in aluminum excephalopathy. Acta Neuropath. 9: 127-133, 1967.
29. Wisniewski, H., and Terry, R.D. Experimental colchicine encephalopathy. I. Induction of neurofibrillary degeneration. Lab. Invest. 17: 577-587, 1967.
30. Terry, R.D., and Wisniewski, H. Neuronal fibrous proteins in some pathological states. Neurosci. Res. Prog. Bull. 6: 184-187, 1968.
31. Rap, Z. M., Wisniewski, H., Werminski, K., and Korthals, J. Thrombosis of the middle cerebral artery in cats induced electrolytically by means of stereotaxis. Neuropat. Pol. 6: 219-226, 1968.
32. Wisniewski, H., Shelanski, M.L., and Terry, R.D. Effects of mitotic spindle inhibitors neurotubules and neurofilaments in anterior horn cells. J. Cell Biol. 38: 224-229, 1968.
33. Wisniewski, H., Weller, R.O., and Terry, R.D. Experimental hydrocephalus produced by the subarachnoid infusion of silicone oil. J. Neurosurg. 31: 10-14, 1969.
34. Bensch, K.G., Marantz, R., Wisniewski, H., and Shelanski, M. Induction in vitro of microtubular crystals by Vinca Alkaloids. Science 165: 495-496, 1969.
35. Shelanski, M.L., and Wisniewski, H. Neurofibrillary degeneration induced by vincristine therapy. A.M.A. Arch. Neurol. 20: 199-206, 1969.

36. Wisniewski, H., Weller, R.O., and Terry, R.D. Experimental hydrocephalus produced by the subarachnoid infusion of silicone oil. J. Neurosurg. 31: 10-14, 1969.
37. Wisniewski, H., Prineas, J., and Raine, C.S. An ultrastructural study of experimental demyelination and remyelination. I. Acute experimental allergic encephalomyelitis in the peripheral nervous system. Lab. Invest. 21: 105-118, 1969.
38. Wisniewski, H., Terry, R.D., Whitaker, J.N., Cook, S.D., and Dowling, P.C. Landry-Guillain-Barre Syndrome. A primary demyelinating disease. A.M.A. Arch. Neurol. 21: 269-276, 1969.
39. Raine, C.S., Wisniewski, H., and Prineas, J. An ultrastructural study of experimental demyelination and remyelination: II. Chronic experimental allergic encephalomyelitis in the central nervous system. Lab. Invest. 21: 316-327, 1969.
40. Prineas, J., Raine, C.S., and Wisniewski, H. An ultrastructural study of experimental demyelination and remyelination: II. Chronic experimental allergic encephalomyelitis in the central nervous system. Lab. Invest. 21: 472-483, 1969.
41. Weller, R.O., Wisniewski, H., Ishii, N., Shulman, K., and Terry, R.D. Brain tissue damage in hydrocephalus. In: Hydrocephalus and Spina Bifida, Developmental Medicine and Child Neurology, Supplement 20, pp. 1-7, 1969.
42. Weller, R.O., and Wisniewski, H. Histological and ultrastructural changes with experimental hydrocephalus in adult rabbits. Brain 92: 819-828, 1969.
43. Raine, C.S., and Wisniewski, H. On the occurrence of microtubules within mature astrocytes. Anat. Record 167: 303-308, 1970.
44. Wisniewski, H., Terry, R.D., and Hirano, A. Neurofibrillary pathology. J. Neuropath. & Exp. Neurol. 29: 163-176, 1970.
45. Wisniewski, H., Johnson, A.B., Raine, C.S., Kay, W.J., and Terry R.D. Senile plaques and cerebral amyloidosis in aged dogs. A histochemical and ultrastructural study. Lab. Invest. 23: 287-296, 1970.
46. Terry, R.D., and Wisniewski, H. The ultrastructure of the neurofibrillary tangle and the senile plaque. In: Alzheimer's Disease and Related Conditions, edited by G.E.W. Wolstenholme, and M. O'Connor. Ciba Foundation Symposium, London, J. & A. Churchill, pp. 145-168, 1970.
47. Wisniewski, H., and Terry, R.D. An experimental approach to the morphogenesis of neurofibrillary degeneration and the argyrophilic plaque. In: Alzheimer's Disease and Related Conditions, edited by G.E.W. Wolstenholme, and M. O'Connor. Ciba Foundation Symposium, London, J. & A. Churchill, pp. 223-243, 1970.
48. Horoupian, D.S., and Wisniewski, H. Neurofilamentous hyperplasia in inferior olivary hypertrophy. J. Neuropath. & Exp. Neurol. 30: 571-582, 1971.
49. Johnson, A.B., Wisniewski, H.M., Raine, C.S., Eylar, E.H., and Terry, R.D. Specific binding of peroxidase-labeled myelin basic protein in allergic encephalomyelitis. Proceedings of the National Academy of Sciences (U.S.A.), 68: 2694-2698, 1971.
50. Weller, R.O., Wisniewski, H., Shulman, K., and Terry, R.D. Experimental hydrocephalus in young dogs. Histological and ultrastructural study of the brain tissue damage. J. Neuropath. & Exp. Neurol. 30: 613-626, 1971.
51. Wisniewski, H.M., and Morell, P. Quaking mouse: Ultrastructural evidence for arrest of myelinogenesis. Brain Res. 29: 63-73, 1971.

52. Raine, C.S., Wisniewski, H.M., Dowling, P.C., and Cook, S.D. An ultrastructural study of experimental demyelination and remyelination. IV. Recurrent episodes and peripheral nervous system plaque formation in experimental allergic encephalomyelitis. Lab. Invest. 25: 28-34, 1971.
53. Wisniewski, H.M., and Raine, C.S. An ultrastructural study of experimental demyelination and remyelination. V. Central and peripheral nervous system lesions caused by Diphtheria toxin. Lab. Invest. 25: 73-80, 1971.
54. Wisniewski, H.M., Coblenz, J.M., and Terry, R.D. Pick's Disease. A clinical and ultrastructural study. A.M.A. Arch. Neurol. 26: 97-108, 1972.
55. Wisniewski, H.M., Raine, C.S., and Kay, W.J. Observations on viral demyelinating encephalomyelitis canine distemper. Lab. Invest. 26: 589-599, 1972.
56. Terry, R.D., and Wisniewski, H.M. Ultrastructure of senile dementia and of experimental analogs. In: Aging and The Brain, edited by C.M. Gaitz. Advances in Behavioral Biology 3: 231, Plenum Press, New York-London, 1972.
57. Ghetti, B., and Wisniewski, H.M. On degeneration of terminals in the cat striate cortex. Brain Res. 44: 630-635, 1972.
58. Ghetti, B., Horoupian, D.S., and Wisniewski, H.M. Transsynaptic response of the lateral geniculate nucleus and the pattern of degeneration of the nerve terminals in the rhesus monkey after eye enucleation. Brain Res. 45: 31-48, 1972.
59. Wisniewski, H.M. Patterns of myelin damage resulting from inflammatory and toxin-induced lesions and their relationship to Multiple Sclerosis. U.C.L.A. Medical Forum. In: Multiple Sclerosis, edited by Wolfram. Academic Press, New York, pp. 53-89, 1972.
60. Morell, P., Greenfield, S., Costantino-Ceccarini, E., and Wisniewski, H. Changes in the protein composition of mouse brain myelin during development. J. Neurochem. 19: 2545-2554, 1972.
61. Wisniewski, H.M., Ghetti, B., and Horoupian, D.S. The fate of synaptic membranes of degenerating optic nerve terminals and their role in the mechanism of transsynaptic changes. J. Neurocytol. 1: 297-310, 1972.
62. Terry, R.D., and Wisniewski, H.M. Pathology of the aging nervous system. Chapter prepared for a survey of Research in the Aging Nervous System, sponsored by the National Institute of Child Health and Development, edited by G. Maletta. National Institutes of Health, Bethesda, 1973.
63. Wisniewski, H.M., and Terry, R.D. Morphology of the aging brain, human and animal. Progress in Brain Res. 40: 184-186, 1973.
64. Wisniewski, H.M., and Terry R.D. Reexamination of the pathogenesis of the senile plaque. In: Progress in Neuropathology, edited by H.M. Zimmerman. Grune & Stratton, New York, Vol 2: 1-26, 1973.
65. Horoupian, D.S., Ghetti, B., and Wisniewski, H.M. Retrograde transneuronal degeneration of optic fibers and their terminals in lateral geniculate nucleus of rhesus monkey. Brain Res. 49: 257-275, 1973.
66. Spencer, P.S., Raine, C.S., and Wisniewski, H. Axon diameter and myelin thickness: Unusual relationships in dorsal root ganglia. Anatom. Record 176: 225-244, 1973.

67. Donat, J.R., and Wisniewski, H.M. The spatiotemporal pattern of Wallerian degeneration in mammalian peripheral nerves. Brain Res. 53: 41-53, 1973.
68. Cook, R.D., and Wisniewski, H.M. The role of oligodendroglia and astroglia in Wallerian degeneration of the optic nerve. Brain Res. 61: 191-206, 1973.
69. Wisniewski, H.M., Ghetti, B., and Terry, R.D. Neuritic (senile) plaques and filamentous changes in aged rhesus monkeys. J. Neuropath. & Exp. Neurol. 32: 566-584, 1973.
70. Morell, P., Greenfield, S., Norton, W.T., and Wisniewski, H. Isolation and characterization of myelin protein from adult quaking mice and its similarity to myelin protein of young normal mice. In: Functional and Structural Proteins of the Nervous System, edited by Davison, Mandel and Morgan. Plenum Publishing Co., New York, pp. 251-261, 1973.
71. Cook, R.D., Raine, C.S., and Wisniewski, H.M. On perivascular astrocytic membrane specializations in monkey optic nerve. Brain Res. 57: 491-497, 1973.
72. Brostoff, S.W., Wisniewski, H.M., Greenfield, S., Morell, P., and Eylar, E.H. Immunopathologic response in guinea pigs sensitized with peripheral nervous system myelin. Brain Res. 58: 500-505, 1973.
73. Coblenz, J.M., Mattis, S., Zingesser, L.H., Kasoff, S.S., Wisniewski, H.M., Katzman, R. Presenile dementia. Clinical aspects and evaluation of cerebrospinal fluid dynamics. Arch. Neurol. 29: 299-308, 1973.
74. Goldfischer, S., Moore, C.L., Johnson, A.B., Spiro, A.J., Valsamis, M.P., Wisniewski, H.M., Ritch, R.H., Norton, W.T., Rapin, I., and Gartner, L.M. Peroxisomal and mitochondrial defects in the cerebro-hepato-renal syndrome. Science 182: 62-64, 1973.
75. Tellez-Nagel, I., and Wisniewski, H.M. Ultrastructure of neurofibrillary tangles in Steele-Richardson-Olszewski syndrome. Arch. Neurol. 29: 324-328, 1973.
76. Brostoff, S.W., Sacks, H., DalCanto, M., Johnson, A.B., Raine, C.S., and Wisniewski, H.M. The P2 protein of bovine root myelin: Isolation and some chemical and immunological properties. J. Neurochem. 23: 1037-1043, 1974.
77. Korthals, J.K., Wisniewski, H.M., Ghetti, B., and Cook, R.D. The fate of the axon and its terminal in the Pacinian corpuscle following sciatic nerve section. J. Neurocytol. 3: 385-403, 1974.
78. Iqbal, K., Wisniewski, H.M., Shelanski, M.L., Brostoff, S., Lwincz, B.H., and Terry, R.D. Protein changes in senile dementia. Brain Res. 77: 337-343, 1974.
79. Kristensson, K., Ghetti, B., and Wisniewski, H.M. Study on the propagation of Herpes Simplex virus (Type 2) into the brain after intraocular injection. Brain Res. 69: 189-201, 1974.
80. Cook, R.D., Ghetti, B., Wisniewski, H.M. The pattern of Wallerian degeneration in the optic nerve of new-born kitten: an ultrastructural study. Brain Res. 75: 261-275, 1974.
81. Dal Canto, M.F., Johnson, A.B., Raine, C.S., Wisniewski, H.M., and Brostoff, S.W. Experimental allergic neuritis: Cells binding horseradish peroxidase conjugates of myelin basic proteins. J. Immunol. 113: 387-394, 1974.
82. Wisniewski, H.M., Brostoff, S.W., Carter, H., and Eylar, E.H. Recurrent episodes of demyelination in experimental allergic polyganglio-radiculo neuritis in rhesus monkey sensitized with rabbit sciatic nerve myelin. Arch. Neurol. 30: 347-358, 1974.

83. Schaumburg, H.H., Wisniewski, H.M., and Spencer, P.S. Ultrastructural studies of dying-back process. 1. Peripheral nerve terminal and axon degeneration in systemic acrylamide intoxication. J. Neuropath. Exper. Neurol. 33: 260-284, 1974.
84. Lhwnicz, B.H., Kristensson, K., Wisniewski, H.M., Shelanski, M.L., and Terry, R.D. Observations on axoplasmic transport in rabbits with aluminum-induced neurofibrillary tangles. Brain Res. 80: 413-420, 1974.
85. Terry, R.D., Wisniewski, H.M. Some structural and chemical aspects of the aging nervous system. Scand. J. Clin. Lab. Invest. 34: 13 (Supplement 141), 1974.
86. Rose, A.L., Wisniewski, H.M., and Cammer, W. Neurotoxicity of hexachlorophene: New pathological and biochemical observations. J. Neurol. Sci. 24: 425, 1975.
87. Wisniewski, H.M., Berry K., and Spiro, A.J. Ultrastructure of thalamic neuronal inclusions in myotonic dystrophy. J. Neurol. Sci. 24: 321-329, 1975.
88. Korthals, J.K., and Wisniewski, H.M. Experimental model of peripheral nerve ischemia. J. Neurol. Sci. 24: 65-76, 1975.
89. Wisniewski, H.M., and Bloom, B.R. Primary demyelination as a non-specific consequence of a cell-mediated immune reaction. J. Experi. Med. 141: 346-359, 1975.
90. Terry, R.D., and Wisniewski, H.M. Pathology and pathogenesis of dementia. In: Neurological and Sensory Disorders in the Elderly, edited by W.S. Fields, pp. 135-150, 1975.
91. Ghetti, B., Horoupian, D.S., and Wisniewski, H.M. Acute and long-term transneuronal response of dendrites of lateral geniculate neurons following transection of the primary visual afferent pathway. In: Advances in Neurology, 12: (Kreutzberg, G.W., ed.) Raven Press, New York, 1975.
92. Carlo, D.J., Karkhanis, Y.D., Bailey, P.J., Wisniewski, H.M., and Brostoff, S.W. Experimental allergic neuritis: Evidence for the involvement of the P₀ and P₂ proteins. Brain Res. 88: 580-584, 1975.
93. Iqbal, K., Wisniewski, H.M., Grundke-Iqbal, I., Korthals, J.K., and Terry, R.D. Chemical pathology of neurofibrils: Neurofibrillary tangles of Alzheimer's presenile-senile dementia. J. Histochem. Cytochem. 23: 563-569, 1975.
94. Wisniewski, H.M., and Bloom, B.R. Experimental allergic optic neuritis (EAON) in the rabbit: A new model to study primary demyelinating disease. J. Neurol. Sci. 24: 313-319, 1975.
95. Wisniewski, H.M., Bruce, M.E., and Fraser, H. Infectious etiology of neuritic (senile) plaques in mice. Science 190: 1108-1110, 1975.
96. Dal Canto, M.C., Wisniewski, H.M., Johnson, A.B., Brostoff, S.W. and Raine, C.S. Vesicular disruption of myelin in autoimmune demyelination. J. Neurol. Sci. 24: 313-319, 1975.
97. Wisniewski, H.M. Morphogenesis of the demyelinating process: Demyelination as a non-specific consequence of a cell-mediated immune reaction. In: Multiple Sclerosis Research, edited by Davison, A.N., Humphrey, J.H., Liversedge, A.L., McDonald, W.I., and Porterfield, J.S., Elsevier, Amsterdam, New York, 132-141, 1975.
98. Hughes, D., Caspary, E.A., and Wisniewski, H.M. Immunosuppression by linoleic acid. Lancet 501, 1975.

99. Berry K., Wisniewski, H.M., Svarzbein, L., and Baez, S. On the relationship of brain vasculature to production of neurological deficit and morphological changes following acute unilateral common carotid artery ligation in gerbils. J. Neurol. Sci. 25: 75-92, 1975.
100. Terry, R.D., and Wisniewski, H.M. Structural and chemical changes of the aged human brain. Aging 2: 127-141, 1975.
101. Wisniewski, H.M., Narang, H.K., and Terry, D. Neurofibrillary tangles of paired helical filaments. J. Neurol. Sci. 27: 173-181, 1976.
102. Iqbal, K., Grundke-Iqbal, I., Wisniewski, H.M., Korthals, J.K., and Terry, R.D. Chemistry of neurofibrillary proteins in aging. In: Neurobiology of Aging, edited by R.D. Terry and S. Gershon, Raven Press, New York, 351-360, 1976.
103. Wisniewski, H.M., and Terry R.D. Neuropathology of the aging brain. In: Neurobiology of Aging, edited by R.D. Terry, and S. Gershon, Raven Press, New York, pp. 265-280, 1976.
104. Brostoff, S.W., Powers, J.M., Wisniewski, H.M., and Hogen, E.L. Encephalitogenic properties of myelin from the quaking mutant. Brain Res. 107: 633-637, 1976.
105. Kristensson, K., Wisniewski, H.M., and Bornstein, M.B. About demyelinating properties of humoral antibodies in experimental allergic encephalomyelitis: in vivo and in vitro studies. Acta Neuropath. (Berl.) 36: 307-314, 1976.
106. Kristensson, K., and Wisniewski, H.M. Penetration of protein tracers into the epiretinal portion of the optic nerve in the rabbit eye. J. Neurol. Sci. 30: 411-416, 1976.
107. Powers, J.M., Balentine, J.D., Wisniewski, H.M., and Terry, R.D. Retroperitoneal ganglioneuroblastoma: A kaleidoscope of neuronal degeneration. A light and electron microscopic study. J. Neuropath. Exp. Neurol. 35: 14-25, 1976.
108. Powers, J.M., Wisniewski, H.M., and Terry, R. Lack of amyloidosis and renal disease in A strain mice. Arch. Pathol. Lab. Med. 100: 69-73, 1976.
109. Wisniewski, H.M. Immunopathology of demyelination in autoimmune diseases and virus infection. Br. Med. Bull. 33: 54-59, 1977.
110. Madrid, R.D., and Wisniewski, H.M. Axonal degeneration in demyelinating disorders. J. Neurocytol. 6: 103, 1977.
111. Smith, A.R., Wilcox, C.B., Eastman, R.E., and Wisniewski, H.M. Cytotoxicity of homogenised lymphocytes in Multiple Sclerosis and other diseases. Lancet I: 138-139, 1977.
112. Raine, C.S., Wisniewski, H.M., Iqbal, K., Grundke-Iqbal, I., and Norton, W.T. Studies on the encephalitogenic effects of purified preparations of human and bovine oligodendrocytes. Brain Res. 120: 269-286, 1977.
113. Narang, H.K., and Wisniewski, H.M. The sequence of myelination in the epiretinal portion of the optic nerve in the rabbit. Neuropath. Appl. Neurobiol. 3: 15-27, 1977.
114. Wisniewski, H.M., and Keith, A.B. Chronic relapsing experimental allergic encephalomyelitis: An experimental model of Multiple Sclerosis. Ann. Neurol. 1: 144-148, 1977.
115. Adams, D.H., Joyce, G., Richardson, V.J., Ryman, B.E., and Wisniewski, H.M. Liposome toxicity in the mouse central nervous system. J. Neurol. Sci. 31: 173-179, 1977.
116. Iqbal, K., Grundke-Iqbal, I., and Wisniewski, H.M. Oligodendroglia from human autopsied brain: Bulk isolation and some chemical properties. J. Neurochem. 28: 707-716, 1977.

117. McDermott, J.R., Iqbal, K., and Wisniewski, H.M. The encephalitogenic activity and myelin basic protein content of isolated oligodendroglia. J. Neurochem. 28: 1081-1088, 1977.
118. Iqbal, K., Wisniewski, H.M., Grundke-Iqbal, I., and Terry, R.D. Neurofibrillary pathology: An update. In: The Aging Brain and Senile Dementia, edited by K. Nandy and I. Sherman. Plenum Press, New York, pp. 209-227, 1977.
119. Stoner, G.L., Brosnan, C.F., Wisniewski, H.M., and Bloom, B.R. Studies on demyelination by activated lymphocytes in the rabbit eye. I. Effects of a mononuclear cell infiltrate induced by products of activated lymphocytes. J. Immunol. 118: 2094-2102, 1977.
120. Brosnan, C.F., Stoner, G.L., Bloom, B.R., and Wisniewski, H.M. Studies on demyelination by activated lymphocytes in the rabbit eye. II. Antibody-dependent cell-mediated demyelination. J. Immunol. 118: 2103-2110, 1977. X
121. Iqbal, K., Grundke-Iqbal, I., Wisniewski, H.M., and Terry, R.D. On neurofilament and neurotubule proteins from human autopsy tissue. J. Neurochem. 29: 417-424, 1977.
122. McDermott, J.R., and Wisniewski, H.M. Studies on the myelin protein changes and antigenic properties of rabbit sciatic nerves undergoing Wallerian degeneration. J. Neurol. Sci. 33: 81-94, 1977.
123. Kristensson, K., and Wisniewski, H.M. Chronic relapsing experimental allergic encephalomyelitis. Studies in vascular permeability changes. Acta Neuropath. (Berl.) 39: 189-194, 1977.
124. McDermott, J.R., Smith, A.I., Iqbal, K., and Wisniewski, H.M. Aluminum and Alzheimer's disease. Lancet II: 710-711, 1977.
125. Wisniewski, H.M., Korthals, J.K., Kopeloff, L.M., Ferszt, R., Chusid, J.C., and Terry, R.D. Neurotoxicity of aluminum. In: Neurotoxicology, edited by L. Roizin and N. Grcevic. Raven Press, New York, pp. 313-315, 1977.
126. Terry, R.D., and Wisniewski, H.M. Structural aspects of aging of the brain. In: Cognitive and Emotional Disturbance in the Elderly, edited by C. Eisdorfer and R. O. Friedel. Yearbook Medical Publishers, Inc., Chicago-London, pp. 3-9, 1977.
127. Iqbal, K., Grundke-Iqbal, I., Terry, R.D., and Wisniewski, H.M. Neurofibrillary proteins. In: Mechanisms, Regulation and Special Function of Protein Synthesis in the Brain, edited by Roberts et al. Elsevier/North Holland Biomedical Press, pp. 171-179, 1977.
128. Iqbal, K., Grundke-Iqbal, I., Wisniewski, H.M., and Terry R.D. Chemical relationship of the paired helical filaments of Alzheimer's dementia to normal human neurofilaments and neurotubules. Brain Res. 142: 321-332, 1978.
129. Kristensson, K. and Wisniewski, H.M. Arrest of myelination and demyelination in rabbit retina induced by herpes simplex virus infection. Neuropath. Applied Neurobiol. 4: 71-82, 1978.
130. Deshmukh, D.S., Bear, W.D., Wisniewski, H.M., and Brockerhoff, H. Long-living liposomes as potential drug carriers. Biochem. Biophys. Res. Commun. 82: 328-334, 1978.
131. Madrid, R.E., and Wisniewski, H.M. Peripheral nervous system pathology in relapsing experimental allergic encephalomyelitis. J. Neurocytol. 7: 265-282, 1978.
132. Loo, Y.H., Scotto, H., and Wisniewski, H.M. Myelin deficiency in experimental phenylketonuria: Contribution of the aromatic acid metabolites of phenylalanine. In: Myelination and Demyelination, edited by J. Palo. Plenum Press, New York and London, pp. 453-469, 1978.

133. Wisniewski, K., Howe, J., Williams, D.G. and Wisniewski, H.M. Precocious aging and dementia in patients with Down's Syndrome. Biol. Psychiatry 13: 619-627, 1978.
134. Lassmann, H., and Wisniewski, H.M. Chronic relapsing EAE - Time course of neurological symptoms and pathology. Acta Neuropathol. (Berl.) 43: 35-42, 1978.
135. Wisniewski, H.M. Possible viral etiology of neurofibrillary changes and neuritic plaques. In: Alzheimer's Disease: Senile Dementia and Related Disorders (Aging Vol. 7), edited by R. Katzman, R.D., Terry and K.L. Bick. Raven Press, New York, pp. 555-558, 1978.
136. Iqbal, K., Grundke-Iqbal, I., Wisniewski, H.M. and Terry, R.D. Neurofibers in Alzheimer dementia and other conditions. In: Alzheimer's Disease: Senile Dementia and Related Disorders (Aging Vol. 7), edited by R. Katzman, R.D., Terry and K.L. Bick. Raven Press, New York, pp. 409-420, 1978.
137. Korthals, J.K., Korthals, M.A. and Wisniewski, H.M. Peripheral nerve ischemia: Part 2. Accumulation of organelles. Annals of Neurol. 4: 487-498, 1978.
138. Wisniewski, H.M. and Soifer, D. Neurofibrillary pathology: Current status and research prospectives. Mechanisms of Ageing and Development 9: 119-142, 1979.
139. Grundke-Iqbal, I., Johnson, A.B., Wisniewski, H.M., Terry, R.D., and Iqbal, K. Evidence that Alzheimer neurofibrillary tangles originate from neurotubules. Lancet i: 578-580, 1979.
140. Wisniewski, K., Jervis, G.A., Moretz, R.C., and Wisniewski, H.M. Alzheimer neurofibrillary tangles in diseases other than senile and presenile dementia. Annals of Neurol. 5: 288-294, 1979.
141. Thormar, H., Wisniewski, H.M., and Lin, F.H. Sera and cerebrospinal fluids from normal uninfected sheep contain a visna virus inhibiting factor. Nature 279: 245-246, 1979.
142. Rose, A.L. and Wisniewski, H.M. Acute bilirubin encephalopathy induced with sulfadimethoxine Gunn rats. J. Neuropath. Exp. Neurol. 38:152-14, 1979.
143. Wisniewski, K., Cobill, J.M., Wilcox, C.B., Caspary, E.A., Gwynn Williams, D. and Wisniewski, H.M. T lymphocytes in patients with Down's syndrome. Biological Psychiatry 14: 463-471, 1979.
144. McDermott, J.R., Smith, A.I., Iqbal, K., and Wisniewski, H.M. Brain aluminum in aging and Alzheimer disease. Neurology 29: 809-814, 1979.
145. Lassmann, H., and Wisniewski, H.M. Chronic relapsing experimental allergic encephalomyelitis: Morphological sequence of myelin degradation. Brain Res. 169: 357-368, 1979.
146. Loo, Y.H., Fulton, T., and Wisniewski, H.M. Vulnerability of the immature brain to phenylacetate intoxication: Tissue permeability to phenylacetate. J. Neurochem. 32: 1697-1698, 1979.
147. Loo, Y.H., Fulton, T., Miller, K.A., and Wisniewski, H.M. Vulnerability of the immature rat brain to phenylacetate intoxication: Postnatal development of the detoxication mechanism. J. Neurochem. 32: 1699-1705, 1979.
148. Madrid, R.E., McDermott, J.R., Pullarkat, R.K., and Wisniewski, H.M. Neuritogenic and chemical properties of guinea pig anterior and posterior root myelin. Brain Research 171: 239-246, 1979.
149. Lassmann, H., and Wisniewski, H.M. Chronic relapsing experimental allergic encephalomyelitis: Effect of age at the time of sensitization on clinical course and pathology. Acta Neuropath. 47: 111-116, 1979.

150. Keith, A.B., Amon, R., Teitelbaum, D., Caspary, E.A., and Wisniewski, H.M. The effect of Cop 1, a synthetic polypeptide, on chronic relapsing experimental allergic encephalomyelitis in guinea pigs. J. Neurol. Sci. 42: 267-274, 1979.
151. Lassmann, H., and Wisniewski, H.M. Chronic relapsing experimental allergic encephalomyelitis. Clinicopathological comparison with Multiple Sclerosis. Arch. Neurol. 36: 490-497, 1979.
152. Mehta, P.D., Kane, A., Thormar, H., and Wisniewski, H.M. Oligoclonal IgG bands and measles antibodies in Multiple Sclerosis (MS) CSF and brain extracts. In: Humoral Immunity In Neurological Diseases, edited by D. Karcher, A. Lowenthal, and A.D. Strosberg. Plenum Publishing Corp., New York, pp. 457-462, 1979.
153. Wisniewski, H.M. Morphology of the aging brain and dementia - human and animal. In: Aging and Immunity, edited by S.K. Singhal, N.R. Sinclair and C.R. Stiller. Elsevier/North Holland, New York/Amsterdam, pp. 185-194, 1979.
154. Wen, G.Y., Sturman, J.A., Wisniewski, H.M., Lidsky, A., Cornwell, A.C., and Hayes, K.C. Tapetum disorganization in taurine-depleted cats. Invest. Ophthalmol. Visual Sci. 18: 1200-1206, 1979.
155. Kristensson, K., Thormar, H., and Wisniewski, H.M. Myelin lesions in the rabbit eye model as a bystander effect of herpes simplex and visna virus sensitization. Acta Neuropath. 48: 215-217, 1979.
156. Grundke-Iqbal, I., Johnson, A.B., Terry, R.D., Wisniewski, H.M., and Iqbal K. Alzheimer neurofibrillary tangles: Antiserum and Immunohistological staining. Ann. Neurol. 6: 532-537, 1979.
157. Wisniewski, H.M. The aging brain. In: Spontaneous Animal Models of Human Disease, edited by E.J. Andrews, B.C. Ward, and N.H. Altman. Academic Press, New York, pp. 148-152, 1979.
158. Czosnek, H.H., Soifer, D., Hochberg, A., and Wisniewski, H.M. Isolation and characterization of free and membrane-bound polyribosomes from rabbit spinal cord. J. Neuroscience Methods 1: 327-341, 1979.
159. Wisniewski, H.M. Neurofibrillary and synaptic pathology in senile dementia of the Alzheimer's type (SDAT). In: Alzheimer's Disease - Early Recognition of Potentially Reversible Deficits, edited by A.I.M. Glen and L.J. Whalley, Churchill Livingstone Edinburgh-London-New York, pp. 7-16, 1979.
160. Haley, J.E., Wisniewski, H.M., and Ledeen, R.W. Extra-axonal diffusion in the rabbit optic system: A caution in myelin and other membranes. Neurochem. Res. 5: 617-628, 1980.
161. Brockerhoff, H., Wisniewski, H.M., Lipton, L.C., and Deshmukh, D.S. Retention of a dialkylphosphatidylcholine in myelin and other membranes. Neurochem. Res. 5: 617-628, 1980.
162. Hetnarski, B., Wisniewski, H.M., Iqbal, K., Dziedzic, J.D., and Lajtha, A. Regional distribution of choline acetyltransferase and acetylcholinesterase in rabbit brain. Neurochem. Res. 5: 385-392, 1980.
163. Brosnan, C.F., and Wisniewski, H.M. Immunopathology of allergic encephalomyelitis. In: Neurochemistry and Clinical Neurology, edited by L. Battistin, G. Hashim, and A. Lajtha. Alan R. Liss, Inc., New York, pp. 379-390, 1980.
164. Wisniewski, K., and Wisniewski, H.M. Diagnosis of infantile neuroaxonal dystrophy by skin biopsy. Ann. Neurol. 7: 377-379, 1980.

165. Czosnek, H., Soifer, D., and Wisniewski, H.M. Heterogeneity of intermediate filament proteins from rabbit spinal cord. Neurochem. Res. 5: 777-793, 1980.
166. Czosnek, H., Soifer, D., and Wisniewski, H.M. Studies on the biosynthesis of neurofilament proteins. J. Cell Biol. 85: 726-734, 1980.
167. Hetnarski, B., Wisniewski, H.M., Iqbal, K., Dziedzic, J.D., and Lajtha, A. Central cholinergic activity in aluminum induced neurofibrillary degeneration. Ann. Neurol. 7: 557-566, 1980.
168. Wen, G.Y., Wisniewski, H.M., Shek, J.W., Loo, Y.H., and Fulton, T.R. Neuropathology of phenylacetate poisoning in rats: An experimental model of phenylketonuria. Ann. Neurol. 7: 557-566, 1980.
169. Wisniewski, H.M., Brosnan, C.F., and Bloom, B.R. Bystander and antibody-dependent cell-mediated demyelination. In: The Suppression of Experimental Allergic Encephalomyelitis and Multiple Sclerosis, edited by A.N. Davison and M.L. Cuzner. Academic Press, London, pp. 45-48, 1980.
170. Hetnarski, B., Lajtha, A., and Wisniewski, H.M. On some derivatives of ferrocene, novel acetylcholinesterase inhibitors. J. Neuroscience Research 5: 1-5, 1980.
171. Wisniewski, H.M., Madrid, R.E., Lassmann, H., Deshmukh, D.S., and Iqbal, K. Search for antigen(s) and immunological mechanisms responsible for extensive demyelination and relapses in experimental allergic encephalomyelitis (EAE). In: Search for the Cause of Multiple Sclerosis and other Chronic Diseases of the Central Nervous System, edited by A. Boese, Verlag Chemie, Weinheim, Germany, pp. 89-95, 1980.
172. Lassmann, H., Kitz, K., and Wisniewski, H.M. Chronic relapsing experimental allergic encephalomyelitis in rats and guinea pigs - A comparison. In: Search for the Cause of Multiple Sclerosis and other Chronic Disease of the Central Nervous System, edited by A. Boese, Verlag Chemie, Weinheim, Germany, pp. 96-104, 1980.
173. Mehta, P.D., Lassmann, H., and Wisniewski, H.M. Immunoglobulin studies in chronic relapsing experimental allergic encephalomyelitis (R-EAE). In: Search for the Cause of Multiple Sclerosis and other Chronic Disorders of the Central Nervous System, edited by A. Boese, Verlag Chemie, Weinheim, Germany, pp. 105-112, 1980.
174. Iqbal, K., Yu, R.K., Grundke-Iqbal, I., and Wisniewski, H.M. Protein and gangliosides of human oligodendroglia. In: Search for the Cause of Multiple Sclerosis and other Chronic Diseases of the Central Nervous System, edited by A. Boese, Verlag Chemie Weinheim, Germany, pp. 139-147, 1980.
175. Iqbal, K., Grundke-Iqbal, I., Johnson, A.B., and Wisniewski, H.M. Neurofibrillary proteins in aging and dementia. In: Aging of the Brain and Dementia (Aging, Vol. 13), edited by L. Amaducci et al. Raven Press, pp. 39-48, 1980.
176. Loo, Y.H., Fulton, T., Miller, K., and Wisniewski, H.M. Phenylacetate and brain dysfunction in experimental phenylketonuria: Synaptic Development. Life Sciences 27: 1283-1290, 1980.
177. Wisniewski, H.M., and Lassmann, H. Chronic relapsing EAE. Its application to the study of human inflammatory demyelinating diseases. In: Progress in Multiple Sclerosis Research, edited by H. Bauer, S. Poser, and G. Ritter, Springer-Verlag, Berlin, pp. 11-17, 1980.
178. Wisniewski, H.M., and Iqbal, K. Ageing of the brain and dementia. Trends in NeuroSciences 3: 226-228, 1980.

179. Mehta, P.D., Thormar, H., and Wisniewski, H.M. Quantitation of measles-specific IgG. Its presence in CSF and brain extracts of patients with multiple sclerosis. Archives of Neurology 37: 607-609, 1980.
180. Grundke-Iqbal, I., Lassmann, H., and Wisniewski, H.M. Chronic relapsing experimental allergic encephalomyelitis. Immunohistochemical studies. Archives of Neurology 37: 651-656, 1980.
181. Wisniewski, H.M., Sturman, J.A., and Shek, J.W. Aluminum chloride induced neurofibrillary changes in the developing rabbit: A chronic animal model. Ann. Neurol. 8: 479-490, 1980.
182. Wisniewski, H.M., Iqbal, K., and McDermott, J.R. Aluminum-induced neurofibrillary changes: Its relationship to senile dementia of the Alzheimer type. Neurotoxicology 1: 121-124, 1980.
183. Lidsky, A.A., Wisniewski, H.M., Madrid, R.E., and Lassmann, H. Visual evoked potentials and pathology in relapsing experimental allergic encephalomyelitis. Docum. Ophthalm. Proc. Series 23: 113-120, 1980.
184. Ramos, P.L., Wisniewski, K., Jervis, G.A., and Wisniewski, H.M. Intermitochondrial septate structures in dystrophic axons. Acta Neuropath. (Berl) 52: 105-109, 1980.
185. Lassmann, H., Kitz, K., and Wisniewski, H.M. Structural variability of demyelinating lesions in different models of subacute and chronic experimental allergic encephalomyelitis. Acta Neuropath. (Berl) 51: 191-201, 1980.
186. Czosnek, H., Soifer, D., Gal, A., Mack, K., Hochberg, A., and Wisniewski, H.M. Poly (A)- and Nonpoly(A)-RNA associated with rat brain microsomal fractions: In vivo labeling studies. J. Neurosci. Res. 5: 515-530, 1980.
187. Thormar, H., Kristensson, K., Lin, F.H., and Wisniewski, H.M. Cellular immune response in rabbits immunized with purified visna virus. Acta Path. Microbiol. Scand. Sect. C. 88: 173-177, 1980.
188. Mehta, P.D., Frisch, S., Thormar, H., Tourtellotte, W.W., and Wisniewski, H.M. Bound antibody in multiple sclerosis brain. J. Neurol. Sci. 49: 91-98, 1981.
189. Robain, O., Wen, G.Y., Wisniewski, H.M., Shek, J.W., and Loo, Y.H. Purkinje cell dendritic development in experimental phenylketonuria. A quantitative analysis. Acta Neuropath. (Berl) 53: 107-112, 1981.
190. Deshmukh, D.S., Kristensson, K., Wisniewski, H.M., and Brockerhoff, H. Toxicity and neuronal transport of stable liposomes and phospholipid in the nervous system. Neurochem. Res. 6: 143-151, 1981.
191. Vorbrodt, A.W., Lossinsky, A.S., Wisniewski, H.M., Moretz, R.C., and Iwanowski, L. Ultrastructural cytochemical studies of cerebral microvasculature in scrapie infected mice. Acta Neuropath. (Berl) 53: 203-211, 1981.
192. Lossinsky, A.S., Vorbrodt, A.W., Wisniewski, H.M. and Iwanowski, L. Ultracytochemical evidence for endothelial channel-lysosome connections in mouse brain following blood-brain barrier changes. Acta Neuropath. (Berl) 53: 197-202, 1981.
193. Czosnek, H., Soifer, D., Mack, K., and Wisniewski, H.M. Similarity of neurofilament proteins from different parts of the rabbit nervous system. Brain Research 216: 387-398, 1981.

194. Schwerer, B., Lassmann, H., Kitz, K., Bernheimer, H., and Wisniewski, H.M. Fractionation of spinal cord tissue affects its activity to induce chronic relapsing encephalomyelitis. Acta Neuropath. Suppl. VII: 165-168, 1981.
195. Lassmann, H., Kitz, K., and Wisniewski, H.M. Ultrastructural variability of demyelinating lesions in experimental allergic encephalomyelitis and multiple sclerosis. Acta Neuropath. Suppl. VII: 173-175, 1981.
196. Kitz, K., Lassmann, H., and Wisniewski, H.M. Isolated leptomeninges of the spinal cord: an ideal tool to study inflammatory reaction in EAE. Acta Neuropath. Suppl. VII: 179-181, 1981.
197. Merz, P.A., Somerville, R.A., Wisniewski, H.M., and Iqbal, K. Abnormal fibrils from scrapie-infected brain. Acta Neuropath. (Berl) 54: 63-74, 1981.
198. Soifer, D., Iqbal, K., Czosnek, H., DeMartini, J., Sturman, J.A., and Wisniewski, H.M. The loss of neuron-specific proteins during the course of Wallerian degeneration of optic and sciatic nerve. J. Neuroscience 1 (5): 461-470, 1981.
199. Madrid, R.E., Wisniewski, H.M., Iqbal, K., Pullarkat, R.K., and Lassmann, H. Relapsing experimental allergic encephalomyelitis induced with isolated myelin and with myelin basic protein plus myelin lipids. J. Neurol. Sci. 50: 399-411, 1981.
200. Lassmann, H., Kitz, K., and Wisniewski, H.M. Histogenesis of demyelinating lesions in the spinal cord of guinea pigs with chronic relapsing experimental allergic encephalomyelitis. J. Neurol. Sci. 50: 109-121, 1981.
201. Moon, H.M., Wisniewski, T., Merz, P., DeMartini, J., and Wisniewski, H.M. Partial purification of neurofilament subunits from bovine brains and studies on neurofilament assembly. J. Cell. Biol. 89: 560-567, 1981.
202. Mehta, P.D., Patrick, B.A., and Wisniewski, H.M. Isoelectric focusing and immunofixation of cerebrospinal fluid and serum in multiple sclerosis. J. Clin. Lab. Immunol. 6: 17-22, 1981.
203. Mehta, P.D., Lassmann, H., and Wisniewski, H.M. Immunologic studies of chronic relapsing EAE in guinea pigs: Similarities to multiple sclerosis. J. Immunology 127: 334-338, 1981.
204. Malik, M.N., Meyers, L.A., Iqbal, K., Sheikh, A.M., Scotto, L., and Wisniewski, H.M. Calcium activated proteolysis of fibrous proteins in central nervous system. Life Sciences 29: 795-802, 1981.
205. Bobin, S.A., Wisniewski, H.M., Kieras, F.J., and Iqbal, K. Morphological and chemical characterization of a starch granule-like polyglucosan deposit isolated from human brain. Acta Neuropath. (Berl) 55: 47-52, 1981.
206. Sturman, J.A., Wen, G.Y., Wisniewski, H.M., and Hayes, K.C. Histochemical localization of zinc in the feline tapetum. Histochemistry 72: 341-350, 1981.
207. Lossinsky, A.S., Vorbrodt, A.W., Wisniewski, H.M., and Moretz, R.C. A simple screening procedure for evaluating central nervous system tissue sections showing structural and cytochemical alterations of the blood-brain barrier. Stain Technology 56: 279-282, 1981.
208. Vorbrodt, A.W., Lassmann, H., Wisniewski, H.M., and Lossinsky, A.S. Ultracytochemical studies of the blood-meningeal barrier (BMB) in rat spinal cord. Acta Neuropath. (Berl) 55: 113-123, 1981.

209. Wisniewski, H.M. Pick disease. In: Handbook of Clinical Neurology, Vol. 42, Neurogenetic Directory, Part 1, edited by P.J. Vinken and G.W. Bruyn. North Holland Publ. Co., Amsterdam, pp. 285-286, 1981.
210. Wisniewski, H.M. Alzheimer disease and senile dementia of the Alzheimer type. In: Handbook of Clinical Neurology, Vol. 42, Neurogenetic Directory, Part 1, edited by P.J. Vinken and G.W. Bruyn. North Holland Publ. Co., Amsterdam, pp. 275-277, 1981.
211. Wisniewski, H.M. Degeneration of the cerebral cortex in infancy, progressive. (Progressive infantile poliodystrophy, Alpers disease). In: Handbook of Clinical Neurology, Vol. 42, Neurogenetic Directory, Part 1, edited by P.J. Vinken and G.W. Bruyn. North Holland Publ. Co., Amsterdam, pp. 486-487, 1981.
212. Lassmann, H., Kitz, K., and Wisniewski, H.M. In vivo effect of sera from animals with chronic relapsing experimental allergic encephalomyelitis on central and peripheral myelin. Acta Neuropath.(Berl) 55: 297-306, 1981.
213. Wisniewski, H.M., Moretz, R.C., and Lossinsky, A.S. Evidence for induction of localized amyloid deposits and neuritic plaques by an infectious agent. Ann. Neurol. 10: 517-522, 1981.
214. Wisniewski, H.M., Sinatra, R.S., Iqbal, K., and Grundke-Iqbal, I. Neurofibrillary and synaptic pathology in the aged brain. In: Aging and Cell Structure, Vol. 1, edited by John E. Johnson, Jr. Plenum Publishing Corp., New York, pp. 105-141, 1981.
215. Lassmann, H., Kitz, K., and Wisniewski, H.M. The development of periventricular lesions in chronic relapsing experimental allergic encephalomyelitis in guinea pigs: A light and scanning electron microscopic study. Neuropathol. Appl. Neurobiol. 7: 1-11, 1981.
216. Dziedzic, J.D., Wisniewski, H.M., and Iqbal, K. Monoamine oxidase activity in normal and Alzheimer brains. Ann. Neurol. 9: 618-619, 1981.
217. Sturman, J.A., Wen, G.Y., Wisniewski, H.M., Niemann, W.H., and Hayes, K.C. Taurine and tapetum structure. In: Taurine in Nutrition and Neurology, edited by Ryan J. Huxtable, and Herminia Pasantes-Morales. Plenum Publishing Corp., New York, pp. 65-78, 1982.
218. Kozlowski, P.B., Moretz, R.C., Carp, R.I., and Wisniewski, H.M. Retinal damage in scrapie mice. Acta Neuropathol. (Berl) 56: 9-12, 1982.
219. Wisniewski, H.M., Lassmann, H., Brosnan, C.F., Mehta, P.D., Lidsky, A.A., and Madrid, R.E. Multiple sclerosis: Immunological and experimental aspects. In: Recent Advances in Clinical Neurology, edited by W.B. Matthews, and G.H. Glaser, No. 3, Churchill Livingstone, Edinburgh, pp. 95-124, 1982.
220. Donadio, M.F., Kozlowski, P.B., Kaplan, H., Wisniewski, H.M., and Majkowski, J. Brain vasculature and induced ischemia in seizure-prone and non-seizure prone gerbils. Brain Research 234: 263-273, 1982.
221. Karcher, D., Lassmann, H., Lowenthal, A., Kitz, K., and Wisniewski, H.M. Antibodies-restricted heterogeneity in serum and cerebrospinal fluid of chronic relapsing experimental allergic encephalomyelitis. J. Neuroimmunology 2: 93-106, 1982.
222. Vorbrodt, A.W., and Wisniewski, H.M. Plasmalemma-bound nucleoside diphosphatase as a cytochemical marker of central nervous system (CNS) mesodermal cells. J. Histochem. Cytochem. 30: 418-424, 1982.
223. Rabe, A., Lee, M.H., Shek, J., and Wisniewski, H.M. Learning deficits in immature rabbits with aluminum-induced neurofibrillary changes. Exp. Neurol. 76: 441-446, 1982.

224. Wisniewski, H.M., Sturman, J.A., and Shek, J.W. Chronic model of neurofibrillary changes induced in mature rabbits by metallic aluminum. Neurobiology of Aging 3: 11-22, 1982.
225. Madrid, R.E., Wisniewski, H.M., Hashim, G.A., Moscarello, M.A., and Wood, D.D. Lipophilin-induced experimental encephalomyelitis in guinea pigs. J. of Neuroscience Research 7: 203-213, 1982.
226. Malik, M.N., Fenko, M.D., Howard, R.G., and Wisniewski, H.M. Further characterization and thiophosphorylation of smooth muscle myosin. Archives of Biochem. & Biophysics 216: 661-670, 1982.
227. Wisniewski, H.M., Vorbrodt, A.W., Moretz, R.C., Lossinsky, A.S., and Grundke-Iqbal, I. Pathogenesis of neuritic (senile) and amyloid plaque formation. In: The Aging Brain - Physiological and Pathophysiological Aspects, edited by S. Hoyer. Experimental Brain Research, Supplementum 5, Springer Verlag, Berlin, Heidelberg, New York, pp. 3-9, 1982.
228. Iqbal, K., Grundke-Iqbal, I., Merz, P.A., and Wisniewski, H.M. Alzheimer neurofibrillary tangle: Morphology and biochemistry. In: The Aging Brain - Physiological and Pathophysiological Aspects, edited by S. Hoyer. Experimental Brain Research, Supplementum 5, Springer Verlag, Berlin, Heidelberg, New York, pp. 10-14, 1982.
229. Grundke-Iqbal, I., Iqbal, K., and Wisniewski, H.M. Immunocytochemical studies on neurofibrillary changes. In: The Aging Brain - Physiological and Pathophysiological Aspects, edited by S. Hoyer. Experimental Brain Research, Supplementum 5, Springer Verlag, Berlin, Heidelberg, New York, pp. 20-25, 1982.
230. Iqbal, K., Grundke-Iqbal, I., Merz, P.A., Wisniewski, H.M. Age-associated neurofibrillary changes. In: The Aging Brain: Cellular and Molecular Mechanisms of Aging in the Nervous System, edited by E. Giacobini, G. Filogamo, G. Giacobini, and A. Vernadakis. Raven Press, New York, pp. 247-257, 1982.
231. Vorbrodt, A.W., Lossinsky, A.S., and Wisniewski, H.M. Cytochemical localization of Quabain-Sensitive, K⁺-dependent p-nitro-phenylphosphatase (transport ATPase) in the mouse central and peripheral nervous system. Brain Research 243: 225-234, 1982.
232. Wen, G.Y., Wisniewski, H.M., and Sturman, J.A. Hereditary abnormality in tapetum lucidum of the Siamese cats. Histochemistry 75: 1-9, 1982.
233. Wisniewski, K., Czosnek, H., Wisniewski, H.M., Soifer, D., Ramos, P.L., Kim, K.S., and Iqbal, K. Reduction of neuronal specific protein and some neurotransmitters in the infantile neuroaxonal dystrophy (INAD). Neuropediatrics 13: 123-129, 1982.
234. Gaull, G.E., Sturman, J.A., Wen, G.Y., and Wisniewski, H.M. Brain and nutrition: The role of taurine. In: Membranes in Growth and Development, edited by J.F. Hoffman, G.H. Giebisch and L. Bolis. Alan R. Liss, Inc., New York, pp. 349-355, 1982.
235. Soifer, D., Czosnek, H.H., Mack, K., Wisniewski, H.M. Properties and dynamics of neurofilament proteins. In: Axoplasmic Transport, edited by D.G. Weiss. Springer-Verlag, Berlin, Heidelberg, New York, pp. 64-72, 1982.
236. Lee, P.K., Deshmukh, D.S., Wisniewski, H.M., and Brockerhoff, H. Axonal transport of phosphatidylcholine and two synthetic analogs. Neurochemistry Intl. 4 (5): 355-359, 1982.
237. Mehta, P.D., Patrick, B.A., Thormar, H., and Wisniewski, H.M. Oligoclonal IgG bands with and without measles antibody activity in sera of patients with subacute sclerosing panencephalitis (SSPE). J. of Immunol. 129: 1983-1985, 1982.

238. Wisniewski, H.M., and Kozlowski, P.B. Evidence for blood-brain barrier changes in senile dementia of the Alzheimer type (SDAT). In: Alzheimer's Disease, Down's Syndrome and Aging, edited by F. Marott Sinex and Carl R. Merrill, Annals of the New York Academy of Sciences, 396: 119-129, 1982.
239. Wisniewski, K.E., French, J.H., Rosen, J.F., Kozlowski, P.B., Tenner, M., and Wisniewski, H.M. Basal ganglia calcification (BGC) in Down's syndrome (DS)—another manifestation of premature aging. In: Alzheimer's Disease, Down's Syndrome and Aging, edited by F. Marrot Sinex and Carl R. Merrill. Annals of the New York Academy of Sciences 396: 179-189, 1982.
240. Wisniewski, H.M., and Loo, Y.H. Phenylketonuria: New clinical and experimental observations. In: Proceedings of the Young Adult Institute 1981 International Conference, edited by J.M. Levy, P.H. Levy, N. Liebman, T.A. Dem, R. Rae, and T.R. Ames. YAI and Workshop Inc., New York, pp. 238-242, 1982.
241. Wen, G.Y., Soifer, D., and Wisniewski, H.M. The doublet microtubules of rods of the rabbit retina. Anat. Embryol. 165: 315-328, 1982.
242. Wen, G.Y., Sturman, J.A., Wisniewski, H.M., MacDonald, A., and Niemann, W.H. Chemical and ultrastructural changes in tapetum of beagles with a hereditary abnormality. Invest. Ophthalmol. Vis. Sci. 23: 733-742, 1982.
243. Wisniewski, H.M., Merz, G.S., Merz, P.A., Wen, G.Y., and Iqbal, K. Morphology and biochemistry of neuronal paired helical filaments and amyloid fibers in humans and animals. In: Progress in Neuropathology, Vol. 5, edited by Harry M. Zimmerman. Raven Press, New York, pp. 139-150, 1983.
244. Moretz, R.C., Wisniewski, H.M., and Lossinsky, A.S. Pathogenesis of neuritic and amyloid plaques in scrapie - ultrastructural study of early changes in the cortical neuropil. In: Aging of the Brain, edited by David Samuel, Sergio Algeri, Samuel Gershon, V.E. Grimm, and Gino Toffano. Raven Press, New York, pp. 61-79, 1983.
245. Brown, W.T., and Wisniewski, H.M. Genetics of Human Aging. Review of Biological Research. Aging 1: 81-99, 1983.
246. Wisniewski, H.M., and Madrid, R.E. Chronic progressive experimental allergic encephalomyelitis (EAE) in adult guinea pigs. J.Neuropath. Exp. Neurol. 42: 243-255, 1983.
247. Malik, M.N., Fenko, M.D., Scotto, L., Merz, P., Rothman, J., Tuzio, H., and Wisniewski, H.M. Purification and characterization of myosin from calf brain. J. Neurochem. 40: 1620-1629, 1983.
248. Lassmann, H., Stemberger, H., Kitz, K., and Wisniewski, H.M. In vivo demyelinating activity of sera from animals with chronic experimental allergic encephalomyelitis. J. Neurol. Sci. 59: 123-137, 1983.
249. Wisniewski, H.M., Brown, H.R., and Thormar, H. Pathogenesis of viral encephalitis: Demonstration of viral antigen(s) in the brain endothelium. Acta Neuropathol (Berl) 60: 107-112, 1983.
250. Merz, P.A., Wisniewski, H.M., Somerville, R.A., Bobin, S.A., Masters, C.L., and Iqbal, K. Ultrastructural morphology of amyloid fibrils from neuritic and amyloid plaques. Acta Neuropathol (Berl) 60: 113-124, 1983.

251. Wisniewski, H.M. Difference in the morphology of Wallerian degeneration in the central nervous system (CNS) and peripheral nervous system (PNS) and its effect on regeneration. In: Nervous System Regeneration, edited by Bernard Haber, J. Regino Perez-Polo, George Hashim, and Anna Maria Gluffrida Stella. Alan R. Liss, New York, March of Dimes Birth Defects Foundation, Birth Defects: Original Article Series, Volume 19, No. 4, pp. 389-395, 1983.
252. Wisniewski, H.M., and Merz, G.S. Neuritic and amyloid plaques in senile dementia of the Alzheimer type. Banbury Report 15: Biological Aspects of Alzheimer's Disease, Cold Spring Harbor, pp. 145-153, 1983.
253. Malik, M.N., Fenko, M.D., Iqbal, K., and Wisniewski, H.M. Purification and characterization of two forms of Ca^{2+} -activated neutral protease from calf brain. J. Biol. Chem. 258: 8955-8962, 1983.
254. Vorbrodt, A.W., Lossinsky, A.S., and Wisniewski, H.M. Enzyme cytochemistry of blood-brain barrier (BBB) disturbances. Acta Neuropathol. (Berl) Suppl. VIII, pp. 43-57, 1983.
255. Wisniewski, H.M., and Lassmann, H. Etiology and pathogenesis of monophasic and relapsing inflammatory demyelination - human and experimental. Acta Neuropathol. (Berl) Suppl. IX, 21-29, 1983.
256. Sturman, J.A., Wisniewski, H.M., and Shek, J.W. High affinity uptake of GABA and glycine by rabbits with aluminum-induced neurofibrillary changes. Neurochem. Res. 8: 1097-1109, 1983.
257. Merz, P.A., Somerville, R.A., and Wisniewski, H.M. Abnormal fibrils in scrapie and senile dementia of the Alzheimer type. In: Virus non Conventionnels et Affections du Systeme Nerveux Central, edited by L.A. Court and F. Cathala. Masson, Paris, pp. 259-281, 1983.
258. Wisniewski, H.M., Merz, G.A., Merz, P.A., and Wen, G.Y. Morphology of the aged brain and senile dementia of the Alzheimer type. In: Neurochemistry of the Aging Brain and Dementia, edited by T. Samorajski and C. Rosten. Texas Research Institute of Mental Sciences, Houston, pp. 43-58, 1983. Presented at the Simposio Internacional "Envejecimiento Cerebral Nuevos Avances" Santiago, Chile, May 13-15, 1982.
259. Thormar, H., Mehta, P.D., Lin, F.H., Brown, H.R., and Wisniewski, H.M. Presence of oligoclonal immunoglobulin G bands and lack of matrix protein antibodies in cerebrospinal fluids and sera of ferrets with measles virus encephalitis. Infection and Immunity 41: 1205-1211, 1983.
260. Robain, O., Wisniewski, H.M., Loo, Y.H., and Wen, G.Y. Experimental phenylketonuria: Effect of phenylacetate intoxication number of synapses in the cerebellar cortex of the rat. Acta Neuropathol. (Berl) 61: 313-315, 1983.
261. Wisniewski, H.M., Lossinsky, A.S., Moretz, R.C., Vorbrodt, A.W., Lassmann, H., and Carp, R.I. Increased blood-brain barrier permeability in scrapie-infected mice. J. Neuropath. Exp. Neurol. 42: 615-626, 1983.
262. Iqbal, K., and Wisniewski, H.M. Neurofibrillary tangles. In: Alzheimer's Disease, edited by R. Reisberg. The Free Press, New York, pp. 48-56, 1983.
263. Wisniewski, H.M. Neuritic (senile) and amyloid plaques. In: Alzheimer's Disease, edited by R. Reisberg. The Free Press, New York, pp. 57-61, 1983.

264. Wisniewski, K.E., and Wisniewski, H.M. Age-associated changes and dementia in Down's syndrome. In: Alzheimer's Disease, edited by B. Reisberg. The Free Press, New York, pp. 319-326, 1983.
265. Wisniewski, H.M., Merz, G.S., Merz, P.A., Wen, G.Y., and Iqbal, K. Neurofibrillary tangles and paired helical filaments in Alzheimer's disease. In: Neurofilaments, edited by C.A. Marotta. University of Minnesota Press, Minneapolis, pp. 196-221, 1983.
266. Lossinsky, A.S., Vorbrodt, A.W., and Wisniewski, H.M. Ultracytochemical studies of vesicular and canicular transport structures in the injured mammalian blood-brain barrier. Acta Neuropathol. (Berl) 61: 239-245, 1983.
267. Merz, G.S., and Wisniewski, H.M. Alzheimer's disease and aging of the human CNS. In: Aging of the Brain, edited by W.H. Gispen and J. Traber. Elsevier Science Publishers B.V., pp. 283-299, 1983.
268. Sturman, J.A., Wen, G.Y., Wisniewski, H.M., and Neuringer, M.D. Retinal degeneration in primates raised on a synthetic human infant formula. Int. J. Devl. Neuroscience 2: 121-129, 1984.
269. Merz, P.A., Somerville, R.A., Wisniewski, H.M., Manuelidis, L., and Manuelidis, E.E. Scrapie-associated fibrils in Creutzfeldt-Jakob disease. Nature 306: 474-476, 1984.
270. Wisniewski, H.M., Loo, Y.H., and Wisniewski, K. Maternal phenylketonuria. Basic research and clinical problems. In: Perspectives and Progress in Mental Retardation, edited by J.M. Berg. Vol. II-Biomedical Aspects, IASSMD, pp. 135-143, 1984.
271. Malik, M.N., Fenko, M.D., and Wisniewski, H.M. Purification and characterization of two forms of Ca^{2+} -activated neutral protease from calf brain synaptosomes and spinal cord. Neurochem. Res. 9: 233-240, 1984.
272. Jenkins, E.C., Brown, W.T., Brooks, J., Duncan, C.J., Rudelli, R.D., and Wisniewski, H.M. Experience with prenatal fragile X detection. Amer. J. of Medical Genetics 17: 215-239, 1984.
273. Iqbal, K., Zaidi, T., Thompson, C.H., Merz, P.A., and Wisniewski, H.M. Alzheimer paired helical filaments: Bulk isolation, solubility, and protein composition. Acta Neuropathol. (Berl) 62: 167-177, 1984.
274. Grundke-Iqbal, I., Iqbal, K., Tung, Y.C., and Wisniewski, H.M. Alzheimer paired helical filaments: Immunochemical identification of polypeptides. Acta Neuropathol. (Berl) 62: 259-267, 1984.
275. Wang, G.P., Grundke-Iqbal, I., Kascsak, R.J., Iqbal, K., and Wisniewski, H.M. Alzheimer neurofibrillary tangles: Monoclonal antibodies to inherent antigen(s). Acta Neuropathol. (Berl) 62: 268-275, 1984.
276. Houthoff, H.J., Moretz, R.C., Rennke, H.G., and Wisniewski, H.M. The role of molecular charge in the extravasation and clearance of protein tracers in blood-brain barrier impairment and cerebral edema. In: Recent Progress in the Study and Therapy of Brain Edema, edited by K.G. Go and A. Baethmann. Plenum Publishing Co., New York, pp. 67-79, 1984.
277. Schuller-Levis, G.B., Clausen, J.T., Schwerer, B., Madrid, R.E., and Wisniewski, H.M. Dynamics of cellular immunity during chronic relapsing EAE in guinea pigs. In: Experimental Allergic Encephalomyelitis: A Useful Model for Multiple Sclerosis, edited by E.C. Alvord, Jr., M.W. Kies, and A.J. Suckling. Alan R. Liss, Inc., New York, pp. 111-117, 1984.

278. Carp, R.I., Merz, G.S., and Wisniewski, H.M. Transmission of unconventional slow virus diseases and the relevance to AD/SDAT transmission studies. In: Senile Dementia: Outlook for the Future, edited by Jean Werthelmer and Maurice Marois. Alan R. Liss, Inc., New York, pp. 31-54, 1984.
279. Mehta, P.D., Patrick, B.A., and Wisniewski, H.M. Oligoclonal IgG band patterns in chronic relapsing EAE and MS. In: Experimental Allergic Encephalomyelitis: A Useful Model for Multiple Sclerosis, edited by E.C. Alvord, Jr., M.W. Kies, and A.J. Suckling. Alan R. Liss, Inc., New York, pp. 341-346, 1984.
280. Wisniewski, H.M., Merz, G.S., and Carp, R.I. Senile dementia of the Alzheimer type: Possibility of infectious etiology in genetically susceptible individuals. Acta Neurologica Scandinavica, Supplementum No. 99, Vol 69, pp. 91-97, 1984.
281. Potempska, A., Loo, Y.H., and Wisniewski, H.M. On the possible mechanism of phenylacetate neurotoxicity: Inhibition of choline acetyltransferase by phenylacetyl-CoA. J. Neurochem. 42: 1499-1501, 1984.
282. Wisniewski, H.M. Neurotoxicity of aluminum. In: Health Protection in Primary Aluminum Production, Vol. 2, Proceedings of a Seminar, Montreal, Sept. 22-24, 1981, edited by James P. Hughes. Intl. Primary Aluminum Institute, London, England, pp. 217-221, 1982. (Distributed 1984).
283. Wisniewski, H.M., Shek, J.W., Gruca, S., and Sturman, J.A. Aluminum-induced neurofibrillary changes in axons and dendrites. Acta Neuropathol. (Berl) 63: 190-197, 1984.
284. Vorbrodt, A.W., Lossinsky, A.S., and Wisniewski, H.M. Ultrastructural studies of concanavalin A receptors and 5'-nucleotidase localization in normal and injured mouse cerebral microvasculature. Acta Neuropathol. (Berl) 63: 210-217, 1984.
285. Gruca, S., and Wisniewski, H.M. Cytochemical study on the effect of aluminum on neuronal Golgi apparatus and lysosomes. Acta Neuropathol. (Berl) 63: 287-295, 1984.
286. Newton, J.C., Barsa-Newton, M.C., Wisniewski, H.M., and Wen, G.Y. Effects of X-radiation on the retina of the albino rabbit as viewed with the transmission electron microscope. Exp. Cell Biology 52: 269-278, 1984.
287. Schwerer, B., Schuller-Levis, G.B., Mehta, P.D., Madrid, R.E., and Wisniewski, H.M. Cellular and humoral immune response to MBP during the course of chronic relapsing EAE. In: Experimental Allergic Encephalomyelitis: A Useful Model for Multiple Sclerosis, edited by E.C. Alvord, Jr., M.W. Kies, and A.J. Suckling. Alan R. Liss, Inc., New York, pp. 187-192, 1984.
288. Vass, K., Lassmann, H., Wisniewski, H.M., and Iqbal, K. Ultracytochemical distribution of myelin basic protein after injection into the cerebrospinal fluid. J. Neurol. Sci. 63: 423-433, 1984.
289. Merz, P.A., Rohwer, R.G., Kascsak, R., Wisniewski, H.M., Somerville, R.A., Gibbs, Jr., C.J., and Gajdusek, D.C. Infection-specific particle from the unconventional slow virus diseases. Science 225: 437-440, 1984.
290. Wisniewski, K.E., Laure-Kamionowska, M., and Wisniewski, H.M. Evidence of arrest of neurogenesis and synaptogenesis in brains of patients with Down's syndrome. New England Journal of Medicine 311: 1187-1188, 1984.
291. Rudelli, R.D., Ambler, M.W., and Wisniewski, H.M. Morphology and distribution of Alzheimer neuritic (senile) and amyloid plaques in striatum and diencephalon. Acta Neuropath. (Berl) 64: 273-281, 1984.

292. Wen, G.Y., and Wisniewski, H.M. Substructures of neurofilaments. Acta Neuropath. (Berl) 64: 339-343, 1984.
293. Wisniewski, H.M., Merz, P.A., and Iqbal, K. Ultrastructure of paired helical filaments of Alzheimer's neurofibrillary tangle. J. Neuropath. Exp. Neurol. 43: 643-656, 1984.
294. Loo, Y.H., Rabe, A., Potempska, A., Wang, P., Fersko, R., and Wisniewski, H.M. Experimental maternal phenylketonuria: An examination of two animal models. Dev. Neurosci. 6: 227-234, 1983/84.
295. Loo, Y.H., Fulton, T.R., Hyde, K.R., and Wisniewski, H.M. Biochemical indices of neuronal development in experimental phenylketonuria: High affinity transport systems and gangliosides. Dev. Neurosci. 6: 235-245, 1983/84.
296. Wisniewski, H.M., Lassmann, H., Schuller-Levis, G., Mehta, P.D., and Madrid, R.E. Pathogenesis of privenous and demyelinating encephalomyelitis and its relevance for multiple sclerosis research. In: Multiple Sclerosis, edited by G. Scarlato and W.B. Matthews, Plenum Publishing Corp., N.Y., pp. 1-21, 1984.
297. Wisniewski, H.M., and Merz, G.S. Neuropathology of the aging brain and dementia of the Alzheimer type. In: Aging 2000: Our Health Care Destiny, edited by C.M. Galtz and T. Samorajski. Volume I, Biomedical Issues, Springer-Verlag, N.Y., pp. 231-243, 1985.
298. Wisniewski, H.M., Schuller-Levis, G.B., Mehta, P.D., Madrid, R.E., and Lassmann, H. Pathogenetic aspects of multiple sclerosis and experimental models of inflammatory demyelination. In: Concepts in Immunopathology, edited by J.M. Cruse and R.E. Lewis, Jr., Vol. 2. S. Karger, Basel, pp. 128-150, 1985.
299. Wisniewski, K.E., Wisniewski, H.M., and Wen, G.Y. Occurrence of neuropathological changes and dementia of Alzheimer's disease in Down's syndrome. Ann. Neurol. 17: 278-282, 1985.
300. Grundke-Iqbal, I., Iqbal, K., Tung, Y.C., Wang, G.P., and Wisniewski, H.M. Alzheimer paired helical filaments: Cross-reacting polypeptide/s normally present in rain. Acta Neuropathol. (Berl) 66: 52-61, 1985.
301. Iqbal, K., Somerville, R.A., Thompson, C.H., and Wisniewski, H.M. Brain glutamate decarboxylase and cholinergic enzyme activities in scrapie. J. Neurol. Sci. 67: 345-350, 1985.
302. Wisniewski, H.M., and Wen, G.Y. Substructures of paired helical filaments from Alzheimer's disease neurofibrillary tangles. Acta Neuropathol. (Berl) 66: 173-176, 1985.
303. Wisniewski, H.M., and Merz, G.S. Aging, Alzheimer's disease and developmental disabilities. In: Aging and Developmental Disabilities: Issues and Approaches, edited by Matthew P. Janicki and Henryk M. Wisniewski, Chapter 10. Brookes Publ. Co., Baltimore, pp. 177-184, 1985.
304. Sturman, J.A., Moretz, R.C., French, J.H., and Wisniewski, H.M. Taurine deficiency in the developing cat: Persistence of the cerebellar external granule cell layer. J. Neurosci. Res. 13: 405-416, 1985.
305. Mehta, P.D., Patrick, B.A., and Wisniewski, H.M. Specificity of oligoclonal IgG bands in sera from chronic relapsing experimental allergic encephalomyelitis guinea pigs. J. Immunology 134: 2338-2342, 1985.

306. Carp, R.I., Merz, P.A., Moretz, R.C., Somerville, R.A., Callahan, S.M., and Wisniewski, H.M. Biological properties of scrapie: An unconventional virus. In: Subviral Pathogens of Plants and Animals: Viroids and Prions, edited by K. Maramorosch and J.J. McKelvey. Academic Press, New York, pp. 425-463, 1985.
307. Sturman, J.A., Moretz, R.C., French, J.H., and Wisniewski, H.M. Taurine deficiency in the developing cat: Persistence of the cerebellar external granule cell layer. Progress in Clin. & Biol. Res. 179: 43-52, 1985.
308. Neuringer, M., Sturman, J.A., Wen, G.Y., and Wisniewski, H.M. Dietary taurine is necessary for normal retinal development in monkeys. Progress in Clin & Biol. Res. 179: 53-62, 1985.
309. Wisniewski, K.E., Dalton, A.J., Crapper McLachlan, D.R., Wen, G.Y., and Wisniewski, H.M. Alzheimer's disease in Down's syndrome: Clinicopathologic studies. Neurology 35: 957-961, 1985.
310. Brown, H.R., Thormar, H., Barshatzky, M., and Wisniewski, H.M. Localization of measles virus antigens in subacute sclerosing panencephalitis in ferrets. Laboratory Animal Science 35: 233-237, 1985.
311. Wisniewski, K., Rudelli, R., Laure-Kamionowska, M., Sklower, S., Houck, Jr., G.E., Kieras, F., Ramos, P., and Wisniewski, H.M. Sanfilippo disease, type A with some features of ceroid lipofuscinosis. Neuropediatrics 16: 98-105, 1985.
312. Sturman, J.A., Moretz, R.C., French, J.H., and Wisniewski, H.M. Postnatal taurine deficiency in the kitten results in a persistence of the cerebellar external granule cell layer: Correction by taurine feeding. J. Neurosci. Res. 13: 521-528, 1985.
313. Wisniewski, H.M., and Wen, G.Y. High-resolution stereo electron microscopy of neurofilaments and Alzheimer type paired helical filaments. In: Proceedings of the 43rd Annual Meeting of the Electron Microscopy Society of America, edited by G.W. Bailey. San Francisco Press, Inc., San Francisco, pp. 730-733, 1985.
314. Mehta, P.D., Thal, L., Wisniewski, H.M., Grundke-Iqbal, I., and Iqbal, K. Paired helical filament antigen in CSF. Lancet 2: 35, 1985.
315. Vorbrodt, A.W., Lossinsky, A.S., Wisniewski, H.M., Suzuki, R., Yamaguchi, T., Masaoka, H., and Klatzo, I. Ultrastructural observations on the transvascular route of protein removal in vasogenic brain edema. Acta Neuropathol. (Berl) 66: 265-273, 1985.
316. Wisniewski, H.M., Merz, G.S., and Carp, R.I. Current hypothesis of the etiology and pathogenesis of senile dementia of the Alzheimer type. Interdiscipl. Topics Geront. 19: 45-53, 1985.
317. Iqbal, K., Grundke-Iqbal, I., and Wisniewski, H.M. Alzheimer neurofibrillary tangles: Biochemical properties. Interdiscipl. Topics Geront. 19: 98-105, 1985.
318. Grundke-Iqbal, I., Iqbal, K., and Wisniewski, H.M. Alzheimer neurofibrillary tangles: Immunochemical cross-reactivity with normal brain antigens. Interdiscipl. Topics Geront. 19: 106-113, 1985.
319. Rudelli, R.D., Brown, W.T., Wisniewski, K., Jenkins, E.C., Laure-Kamionowska, M., Connell, F., and Wisniewski, H.M. Adult Fragile X syndrome. Clinico-neuropathologic findings. Acta Neuropathol. (Berl) 67: 289-295, 1985.
320. Carp, R.I., Merz, P.A., Kascsak, R.J., Merz, G.S., and Wisniewski, H.M. Nature of the scrapie agent: Current status of facts and hypothesis. J. Gen. Virol. 66: 1357-1368, 1985.

321. Kascsak, R.J., Rubenstein, R., Merz, P.A., Carp, R.I., Wisniewski, H.M. and Diringer, H. Biochemical differences and scrapie-associated fibrils support the biological diversity of scrapie agents. J. Gen. Virol. 66: 1715-1722, 1985.
322. Wisniewski, H.M. Progressive supranuclear palsy. The Steele-Richardson-Olszewski syndrome. In: Handbook of Clinical Neurology, edited by P.J. Vinken, G.W. Bruyn, H.L. Klawans, Elsevier Science Publishers, Amsterdam, The Netherlands, 1985, pp. 301-303.
323. Wisniewski, K., Sturman, J.A., Devine, E., Brown, W.T., Rudelli, R., and Wisniewski, H.M. Cystathionine disappearance with neuronal loss: A possible neuronal marker. Neuropediatrics 16: 126-130, 1985.
324. Merz, P.A., Kascsak, R., Rubenstein, R., Carp, R.I., and Wisniewski, H.M. Variations in SAF from different scrapie agents. In: Proceedings of Workshop on Slow Transmissible Diseases, edited by J. Tateishi, pp. 137-145, 1985. Research Committee on Slow Virus Infection. The Japanese Ministry of Health and Welfare.
325. Mehta, P.D., Patrick, B.A., Mehta, S.P., and Wisniewski, H.M. Chronic relapsing EAE in guinea pigs: IgG index and oligoclonal bands in cerebrospinal fluid and sera. Immunol. Investigations 14: 347-354, 1985.
326. Wisniewski, H.M., Merz, G.S., Wen, G.Y., Iqbal, K., and Grundke-Iqbal, I. Morphology and biochemistry of Alzheimer's disease. In: Senile Dementia of the Alzheimer Type, edited by J.T. Hutton and A.D. Kenny, Alan R. Liss, Inc., New York, 1985, pp. 263-274.
327. Wisniewski, H.M., and Wen, G.Y. A comparative study on the substructures of neurofilaments and paired helical filaments from Alzheimer neurofibrillary tangles. Annals N.Y. Academy Sci. 455: 814-815, 1985.
328. Loo, Y.H., Hyde, K.R., Lin, F.H., and Wisniewski, H.M. Cerebral biochemical abnormalities in experimental maternal phenylketonuria: Gangliosides and sialoglycoproteins. Life Sciences 37: 2099-2109, 1985.
329. Loo, Y.H., Potempska, A., and Wisniewski, H.M. A biochemical explanation of phenyl acetate neurotoxicity in experimental phenylketonuria. J. Neurochem. 45: 1596-1600, 1985.
330. Wen, G.Y., and Wisniewski, H.M. Histochemical localization of aluminum in the rabbit CNS. Acta Neuropathol. (Berl) 68: 175-184, 1985.
331. Grundke-Iqbal, I., Wang, G.P., Iqbal, K., Tung, Y.C., and Wisniewski, H.M. Alzheimer paired helical filaments: Identification of polypeptides with monoclonal antibodies. Acta Neuropathol. (Berl) 68: 279-283, 1985.
332. Wisniewski, H.M., Sturman, J.A., Shek, J.W., and Iqbal, K. Aluminum and the central nervous system. J. Environmental Pathol. Toxicol. & Oncol. 6: 1-8, 1985.
333. Sanz, M.M., Jenkins, E.C., Brown, W.T., Davisson, M.T., Kevin, M.J., Roderick, T.H., Silverman, W.P., and Wisniewski, H.M. Mouse chromosome fragility. Am. J. Med. Genetics 23: 491-509, 1986.
334. Vorbrodt, A.W., Dobrogowska, D.H., Lossinsky, A.S., and Wisniewski, H.M. Ultrastructural localization of lectin receptors on the luminal and abluminal aspects of brain. J. Histochem. Cytochem. 34: 251-261, 1986.
335. Lassmann, H., Vass, K., Brunner, C., and Wisniewski, H.M. Peripheral nervous system lesions in experimental allergic encephalomyelitis. Acta Neuropathol. (Berl) 69: 193-204, 1986.

350. Shek, J.W., Wen, G.Y., and Wisniewski, H.M. Atlas of the Rabbit Brain and Spinal Cord, Karger, Basel-New York, 139 pages, 1986.
351. Kim, Y.S., Lee, M.H., and Wisniewski, H.M. Aluminum induced reversible change in permeability of the blood-brain barrier to [14C] sucrose. Brain Res. 377: 286-291, 1986.
352. Vorbrodt, A.W., Lossinsky, A.S., Dobrogowska, D.H., and Wisniewski, H.M. Distribution of anionic sites and glycoconjugates on the endothelial surfaces of the developing blood-brain barrier. Dev. Brain Res. 29: 69-79, 1986.
353. Malik, M.N., Sheikh, A.M., Fenko, M.D., and Wisniewski, H.M. Purification and degradation of purified neurofilament proteins by the brain calcium-activated neutral proteases. Life Sci. 39: 1335-1343, 1986.
354. Iqbal, K., Grundke-Iqbal, I., Zaidi, T., Merz, P.A., Wen, G.Y., Shalkh, S.S., Wisniewski, H.M., Alafuzoff, I., and Winblad, B. Defective brain microtubule assembly in Alzheimer's disease. Lancet 2: 421-426, 1986.
355. Kascsak, R.J., Rubenstein, R., Merz, P.A., Carp, R.I., Robakis, N.K., Wisniewski, H.M., and Diringer, H. Immunological comparison of scrapie-associated fibrils isolated from animals infected with four different scrapie strains. J. Virol. 59: 676-683, 1986.
356. Merz, P.A., Wisniewski, H.M., Rubenstein, R., and Kascsak, R.J. Immunological studies on paired helical filaments and amyloid of Alzheimer's disease. In: Discussions in Neurosciences, edited by A. Bignami, L. Bolis, and D.C. Gajdusek, Volume III, FESN, pp. 58-68, 1986.
357. Merz, P.A., Kascsak, R.J., Robakis, N., Rubenstein, R., and Wisniewski, H.M. Recent studies of scrapie associated fibrils (SAF). In: Discussions in Neurosciences, edited by A. Bignami, L. Bolis, and D.C. Gajdusek, Volume III, FESN, pp. 91-94, 1986.
358. Wisniewski, H.M., Merz, G.S., Merz, P.A., Wen, G.Y., Iqbal, K., Grundke-Iqbal, I., Bobin, S.A., Currie, J., and Miller, D. Abnormal fibers, Alzheimer's disease and unconventional slow infections. In: Senile Dementias: Early Detection, edited by A. Bes, J. Cahn, R. Cahn, S. Hoyer, J.P. Marc-Vergnes and H.M. Wisniewski, Current Problems in Senile Dementias, No. 1. John Libbey Eurotext, London-Paris, pp. 202-210, 1986.
359. Imaki, H., Moretz, R.C., Wisniewski, H.M., and Sturman, J.A. Feline maternal taurine deficiency: Effects on retina and tapetum of the offspring. Dev. Neurosci. 8: 160-181, 1986.
360. Wisniewski, H.M., Wen, G.Y., Wang, K.C., Iqbal, K., and Rubenstein, R. Determination of the handedness of paired helical filaments in Alzheimer's disease. In: Electron Microscopy and Alzheimer's Disease, edited by J. Metuzals. San Francisco Press, Inc., California, pp. 21-24, 1986.
361. Wisniewski, H.M., Iqbal, K., Grundke-Iqbal, I., Rubenstein, R., Wen, G.Y., Merz, P.A., Kascsak, R., and Kristensson, K. Amyloid in Alzheimer's disease and unconventional viral infections. International Symposium on Dementia and Amyloid, "Neuropathology" Supplement 3, The Japanese Soc. of Neuropath., Sasaki Printing & Publishing Co., Ltd, Japan, pp. 87-94, 1986.
362. Wisniewski, H.M., Moretz, R.C., and Iqbal, K. No evidence for aluminum in the etiology and pathogenesis of Alzheimer's disease. Neurobiology of Aging 7: 532-535, 1986.
363. Wisniewski, H.M., Rudelli, R.D., Iqbal, K., and Merz, G. AD/SDAT, plaques, tangles and BBB changes. Neurobiology of Aging 7: 504, 1986.

364. Wisniewski, H.M. and Miller, D.L. Morphology and biochemistry of abnormal fibrous proteins in Alzheimer's disease and senile dementia of the Alzheimer type (SDAT). Neurobiology of Aging 7: 446-447, 1986.
365. Wisniewski, H.M., and Rabe, A. Discrepancy between Alzheimer-type neuropathology and dementia in persons with Down's syndrome. In: Mental Retardation: Research, Education, and Technology Transfer, edited by Henryk M. Wisniewski and Donald A. Snider, The New York Academy of Sciences, New York, NY. Annals N.Y. Acad. Sci. 477: 247-260, 1986.
366. Iqbal, K., Grundke-Iqbal, I., and Wisniewski, H.M. Neuronal cytoskeleton in aging and dementia. In: Progress in Brain Research, edited by D.F. Swaab, E. Fliers, M. Mirmiran, W.A. Van Gool and F. Van Haaren. Elsevier Science Publishers (Biomedical Division), Amsterdam, The Netherlands, pp. 279-288, 1986.
367. Crapper McLachlan, D.R., Lukiw, W.J., Cho, H.J., Carp, R.I., and Wisniewski, H.M. Chromatin structure in scrapie and Alzheimer's disease. Can. J. Neurol. Sci. 13: 427-431, 1986.
368. Iqbal, K., Grundke-Iqbal, I., and Wisniewski, H.M. Alzheimer's disease, microtubule and neurofilament proteins, and axoplasmic flow. Lancet 1: 102, 1987.
369. Malik, M.N., Ramaswamy, S., Tuzio, H., Shiekh, A.M., Fenko, M.D., Wisniewski, H.M., and Howard, R.G. Micromolar Ca^{2+} requiring protease from human platelets: purification, partial characterization and effect on the cytoskeletal proteins. Life Sci. 40: 593-604, 1987.
370. Robakis, N.K., Wisniewski, H.M., Jenkins, E.C., Devine-Gage, E.A., Houck, G.E., Yao, X-L., Ramakrishna, N., Wolfe, G., Silverman, W. P., and Brown, W.T. Chromosome 21q21 sublocalisation of gene encoding beta-amyloid peptide in cerebral vessels and neuritic (senile) plaques of people with Alzheimer disease and Down syndrome. Lancet 1: 384-385, 1987.
371. Merz, G.S., Schwenk, V., Schuller-Levis, G., Gruca, S., and Wisniewski, H.M. Isolation and characterization of macrophages from scrapie-infected mouse brain. Acta Neuropathol. (Berl) 72: 240-247, 1987.
372. Merz, P.A., Kascsak, R.J., Rubenstein, R., Carp, R.I., and Wisniewski, H.M. Antisera to scrapie-associated fibril protein and prion protein decorate scrapie-associated fibrils. J. Virol. 61: 42-49, 1987.
373. Cook, R.D., and Wisniewski, H.M. The spatio-temporal pattern of Wallerian degeneration in the rhesus monkey optic nerve. Acta Neuropathol. (Berl) 72: 261-267, 1987.
374. Mehta, P.D., Patrick, B.A., Mehta, S.P., and Wisniewski, H.M. Specificity of oligoclonal IgG bands against myelin proteins in chronic relapsing eae in guinea pigs. J. Immunol. 138: 746-751, 1987.
375. Wen, G.Y., and Wisniewski, H.M. High resolution analysis of paired helical filaments in Alzheimer's disease. J. Electron Microscopy Technique 5: 347-355, 1987.
376. Wu, Y., Brown, W.T., Robakis, N.K., Dobkin, C., Devine-Gage, E., Merz, P., and Wisniewski, H.M. A pvuii rfp detected in the human prion protein (prp) gene. Nucleic Acids Research 15: 3191, 1987.
377. Szumanska, G., Vorbrodt, A.W., Mandybur, T.I., and Wisniewski, H.M. Lectin histochemistry of plaques and tangles in Alzheimer's disease. Acta Neuropathol. (Berl) 73: 1-11, 1987.

336. Szumanska, G., Vorbrodt, A.W., and Wisniewski, H.M. Lectin histochemistry of scrapie amyloid plaques. Acta Neuropathol. (Berl) 69: 205-212, 1986.
337. Palackal, T., Moretz, R., Wisniewski, H., and Sturman, J. Abnormal visual cortex development in the kitten associated with maternal dietary taurine deprivation. J. Neurosci. Res. 15: 223-239, 1986.
338. Iqbal, K., Grundke-Iqbal, I., Zaidi, T., Ali, N., and Wisniewski, H.M. Are Alzheimer neurofibrillary tangles insoluble polymers? Life Sciences 38: 1695-1700, 1986.
339. Grundke-Iqbal, I., Iqbal, K., Quinlan, M., Tung, Y-C., Zaidi, M.S., and Wisniewski, H.M. Microtubule-associated protein Tau. J. Biol. Chem. 261: 6084-6089, 1986.
340. Lossinsky, A.S., and Wisniewski, H.M. A comparative ultrastructural study of endothelial cell tubular structures from injured mouse blood-brain barrier and normal hepatic sinusoids demonstrated after perfusion fixation with osmium tetroxide. Microvascular Research 31: 333-344, 1986.
341. Wisniewski, H.M. Clinical, pathological and biochemical aspects of Alzheimer's disease. In: Alzheimer's and Parkinson's Diseases, edited by A. Fisher, I. Hanin and C. Lachman. Plenum Publishing Corp., New York, pp. 25-36, 1986.
342. Iqbal, K., Grundke-Iqbal, I., and Wisniewski, H.M. Alzheimer neurofibrillary tangle and its relationship with plaque core amyloid. In: Amyloidosis, edited by G.G. Glenner, E.F. Osseman, E.P. Benditt, E. Calkins, A.S. Cohen, and D. Zucker-Franklin. Plenum Publishing Corp., New York, pp. 717-722, 1986.
343. Rubenstein, R., Kascsak, R.J., Merz, P.A., Wisniewski, H.M., Carp, R.I., and Iqbal, K. Paired helical filaments associated with Alzheimer disease are readily soluble structures. Brain Res. 372: 80-88, 1986.
344. Rubenstein, R., Kascsak, R.J., Merz, P.A., Papini, M.C., Carp, R.I., Robakis, N.K., and Wisniewski, H.M. Detection of scrapie-associated fibril (SAF) proteins using anti-SAF antibody in non-purified tissue preparations. J. Gen. Virol. 67: 671-681, 1986.
345. Vass, K., Lassmann, H., Wekerle, H., and Wisniewski, H.M. The distribution of Ia antigen in the lesions of rat acute experimental allergic encephalomyelitis. Acta Neuropathol. (Berl) 70: 149-160, 1986.
346. Schuller-Levis, G.B., Kozlowski, P.B., and Wisniewski, H.M. Cyclosporin A treatment of an induced attack in a chronic relapsing model of experimental allergic encephalomyelitis. Clin. Immun. & Immunopath. 40: 244-252, 1986.
347. Grundke-Iqbal, I., Iqbal, K., Tung, Y-C., Quinlan, M., Wisniewski, H.M., and Binder, L.I. Abnormal phosphorylation of the microtubule-associated protein (tau) in Alzheimer cytoskeletal pathology. Proc. Natl. Acad. Sci. USA 83: 4913-4917, 1986.
348. Vorbrodt, A.W., Lossinsky, A.S., and Wisniewski, H.M. Localization of alkaline phosphatase activity in endothelia of developing and mature mouse blood-brain barrier. Dev. Neurosci. 8: 1-13, 1986.
349. Lossinsky, A.S., Vorbrodt, A.W., and Wisniewski, H.M. Characterization of endothelial cell transport in the developing mouse blood-brain barrier. Dev. Neurosci. 8: 61-75, 1986.

378. Lossinsky, A.S., Moretz, R.C., Carp, R.I., and Wisniewski, H.M. Ultrastructural observations of spinal cord lesions and blood-brain barrier changes in scrapie-infected mice. Acta Neuropathol. (Berl) 73: 43-52, 1987.
379. Kim, Y.S., Carp, R.I., Callahan, S.M., and Wisniewski, H.M. Incubation periods and survival times for mice injected stereotactically with three scrapie strains in different brain regions. J. Gen. Virol. 68: 695-702, 1987.
380. Wisniewski, H.M., Iqbal, K., Grundke-Iqbal, I., and Rubenstein, R. The solubility controversy of paired helical filaments: A commentary. Neurochemical Res. 12: 93-95, 1987.
381. Kim, Y.S., Carp, R.I., Callahan, S.M., and Wisniewski, H.M. Scrapie-induced obesity in mice. J. Infect. Diseases 156: 402-405, 1987.
382. Robakis, N.K., Ramakrishna, N., Wolfe, G., and Wisniewski, H.M. Molecular cloning and characterization of a cDNA encoding the cerebrovascular and neuritic plaque amyloid peptides. Proc. Natl. Acad. Sci. (USA) 84: 4190-4194, 1987.
383. Bancher, C., Lassmann, H., Budka, H., Grundke-Iqbal, I., Iqbal, K., Wiche, G., Seitelberger, F., and Wisniewski, H.M. Neurofibrillary tangles in Alzheimer's disease and progressive supranuclear palsy: antigenic similarities and differences. Acta Neuropathol. (Berl) 74: 39-46, 1987.
384. Rubenstein, R., Merz, P.A., Kascsak, R.J., Carp, R.I., Scalici, C.L., Fama, C.L., and Wisniewski, H.M. Detection of scrapie-associated fibrils (SAF) and SAF proteins from scrapie-affected sheep. J. Infect. Disease 156: 36-42, 1987.
385. Popovitch, E.R., Wisniewski, H.M., Kaufman, M.A., Grundke-Iqbal, I., and Wen, G.Y. Young adult-form of dementia with neurofibrillary changes and Lewy bodies. Acta Neuropathol. (Berl) 74: 97-104, 1987.
386. Wisniewski, H.M., and Wen, G.Y. Neurofibrillary changes. In: Encyclopedia of Neuroscience, edited by George Adelman. Birkhauser, Boston, Volume II, pp. 783-786, 1987.
387. Wisniewski, H.M., Wen, G.Y., Grundke-Iqbal, I., Iqbal, K., Mehta, P.D., Merz, P.A., Miller, D., Currie, J., Bobin, S.A., Robakis, N.K., Rabe, A., Brown, W.T. Pathological, biochemical, and genetic aspects of Alzheimer's disease in aged people and Down syndrome. In: Contributions of Chemistry to Health, edited by H. Machleidt. Proceedings of the Fifth CHEMRAWN Conference, Heidelberg, 1986, pp. 297-315, 1987.
388. Iqbal, K., Grundke-Iqbal, I., Merz, P.A., Wisniewski, H.M., and Zaidi, T. *In vitro* assembly and isolation of neurofilaments and microtubules from mammalian CNS. Molecular Brain Research 2: 163-172, 1987.
389. Kozlowski, P.B., Schuller-Levis, G.B., and Wisniewski, H.M. Induction of synchronized relapses in SJL/J mice with chronic relapsing experimental allergic encephalomyelitis. Acta Neuropathol. (Berl) 74: 163-168, 1987.
390. Malik, M.N., Fenko, M.D., Sheikh, A.M., Kascsak, R.J., Tonna-DeMasi, M.S., and Wisniewski, H.M. Third form of calcium-activated neutral proteinase from calf brain: purification, partial characterization and comparison of properties with other forms. Biochim. Biophys. Acta 916: 135-144, 1987.
391. Reeves, R.H., Robakis, N.K., Oster-Granite, M.L., Wisniewski, H.M., Coyle, J.T., and Gearhart, J.D. Genetic linkage in the mouse of genes involved in Down syndrome and Alzheimer's disease in man. Molecular Brain Research 2: 215-221, 1987.

392. Bobin, S.A., Currie, J.R., Merz, P.A., Miller, D.L., Styles, J., Walker, W.A., Wen, G.Y., and Wisniewski, H.M. The comparative immunoreactivities of brain amyloids in Alzheimer's disease and scrapie. Acta Neuropathol. (Berl) 74: 313-323, 1987.
393. Jenkins, E.C., Devine-Gage, E.A., Yao, X-L., Houck, Jr., G.E., Brown, W.T., Wisniewski, H.M., and Robakis, N.K. In-situ hybridisation of the beta-amyloid protein probe to chromosome 9 in patients with familial Alzheimer's disease. Lancet 2: 1155-1156, 1987.
394. Wisniewski, H.M., Bendheim, P.E., and Bolton, D.C. Alzheimer's disease, Down syndrome, and Parkinson's disease: Selected pathologic features. In: Alzheimers Disease, edited by H.J. Altman, Plenum Publishing Corp., New York, pp. 87-97, 1987.
395. Sturman, J.A., Palackal, T., Imaki, H., Moretz, R.C., French, J., and Wisniewski, H.M. Nutritional taurine deficiency and feline pregnancy and outcome. In: The Biology of Taurine, edited by R.J. Huxtable, F. Franconi and A. Giotti. Plenum Publishing Corp., New York, pp. 113-124, 1987.
396. Malik, M.N., Fenko, M.D., Sheikh, A.M., Kasczak, R.J., Tonna-DeMasi, M.S., and Wisniewski, H.M. Third form of calcium-activated neutral proteinase from calf brain: purification, partial characterization and comparison of properties with other forms. Biochim. et Biophys. Acta 916: 135-144, 1987.
397. Wisniewski, K.E., Rabe, A., and Wisniewski, H.M. Commentary on: Pathological similarities between Alzheimer's disease and Down's syndrome: Is there a genetic link? by M.J. Ball, Integr. Psychiatry 5: 159-170, 1987.
398. Brown, H.R., Goller, N.L., Thormar, H., Rudelli, R., Tourtellotte, W.W., Shapshak, P., Boostanfar, R., and Wisniewski, H.M. Measles virus matrix protein gene expression in a subacute sclerosing panencephalitis patient brain and virus isolate demonstrated by cDNA hybridization and immunocytochemistry. Acta Neuropathol. (Berl) 75: 123-130, 1987.
399. Lee, M.H., Rabe, A., Currie, J.R., Shek, J., and Wisniewski, H.M. Transplants of normal fetal cerebral cortical tissue into congenitally malformed brains of infant rats. Ann. N.Y. Acad. Sci. 495: 732-735, 1987.
400. Kasczak, R.J., Rubenstein, R., Merz, P.A., Tonna-DeMasi, M., Fersko, R., Carp, R.I., Wisniewski, H.M. Mouse polyclonal and monoclonal antibody to scrapie-associated fibril proteins. J. Virol. 61: 3688-3693, 1987.
401. Robakis, N.K., Wolfe, G., Ramakrishna, N., and Wisniewski, H.M. Isolation of a cDNA clone encoding the Alzheimer's disease and Down's syndrome amyloid peptide. Banbury Report 27: Molecular Neuropathology of Aging, Cold Spring Harbor Laboratory, pp. 267-281, 1987.
402. Wisniewski, H.M., Rabe, A., and Wisniewski, K.E. Neuropathology and dementia in people with Down's syndrome. Banbury Report 27: Molecular Neuropathology of Aging, Cold Spring Harbor Laboratory, pp. 399-413, 1987.
403. Iqbal, K., Grundke-Iqbal, I., and Wisniewski, H.M. Alterations in the neuronal cytoskeleton in Alzheimer disease. In: Alterations in the Neuronal Cytoskeleton in Alzheimer Disease, edited by G. Perry. Plenum Publishing Corp, pp. 109-136, 1987.
404. Imaki, H., Moretz, R., Wisniewski, H.M., Neuringer, M., and Sturman, J. Retinal degeneration in 3-month-old rhesus monkey infants fed a taurine-free human infant formula. J. of Neuroscience Res. 18: Alan R. Liss, pp. 602-614, 1987.

405. Vorbrodt, A.W., Dobrogowska, D.H., Kim, Y.S., Lossinsky, A.S., and Wisniewski, H.M. Ultrastructural studies of glycoconjugates in brain micro-blood vessels and amyloid plaques of scrapie-infected mice. Acta Neuropathol. (Berl) 75: 277-287, 1988.
406. Jenkins, E.C., Devine-Gage, E.A., Robakis, N.K., Yao, X-L., Brown, W.T., Houck, Jr., G.E., Wolfe, G., Ramakrishna, N., Silverman, W.P., and Wisniewski, H.M. Fine mapping of an Alzheimer disease-associated gene encoding beta-amyloid protein. Biochem. Biophys. Res. Comm., 151: 1-8, 1988.
407. Wisniewski, H.M., and Wrzolek, M. Pathogenesis of amyloid formation in Alzheimer's disease, Down's syndrome and scrapie. In: Novel Infectious Agents and the Central Nervous System, Ciba Foundation Symposium 135, John Wiley & Sons, Chichester, pp. 224-238, 1988.
408. Wisniewski, H.M., Currie, J.R., Barcikowska, M., Robakis, N.K., and Miller, D.L. Alzheimer's disease, a cerebral form of amyloidosis. In: Immunology and Alzheimer's Disease, edited by A. Pouplard-Barthelax, J. Emile and Y. Christen. Springer-Verlag, Paris, pp. 1-6, 1988.
409. Palackal, T., Moretz, R.C., Wisniewski, H.M., and Sturman, J.A. Ultrastructural abnormalities in the visual cortex of kittens from taurine-deficient mothers. Brain Dysfunction 1: 71-89, 1988.
410. Kim, K.S., Miller, D.L., Sapienza, V.J., Chen, C.M.J., Chun, B., Grundke-Iqbal, I., Currie, J.R., and Wisniewski, H.M. Production and characterization of monoclonal antibodies reactive to synthetic cerebrovascular amyloid peptide. Neurosci. Res. Commun. 2: 121-130, 1988.
411. Soifer, D., Mack, K., Wisniewski, H.M. A cDNA coding for rabbit neurofilament protein H: A window on the role of H in neurodegenerative diseases. In: Neural Development and Regeneration, NATO ASI Series, Vol. H22, edited by A. Gorio, J.R. Perez-Polo, J. de Vellis and B. Haber. Springer Verlag, Heidelberg, pp. 333-341, 1988.
412. Loo, Y.H., Wisniewski, K.E., Hyde, K.R., Fulton, T.R., Lin, Y.Y., Wisniewski, H.M. The neurotoxic metabolite of phenylalanine in phenylketonuria. In: Dietary Phenylalanine and Brain Function, edited by R.J. Wurtman and E. Ritter-Walker. Birkhauser Boston - Basel, pp. 249-253, 1988.
413. Hart, M.N., Merz, P., Bennett-Gray, J., Menezes, A.H., Goeken, J.A., Schelper, R.L., Wisniewski, H.M. Beta-amyloid protein of Alzheimer's disease is found in cerebral and spinal cord vascular malformations. Amer. Journal of Pathol. 132: 167-172, 1988.
414. Grundke-Iqbal, I., Vorbrodt, A.W., Iqbal, K., Tung, Y.C., Wang, G.P., Wisniewski, H.M. Microtubule-associated polypeptides tau are altered in Alzheimer paired helical filaments. Molecular Brain Res. 4: 43-52, 1988.
415. Wisniewski, H.M., Wen, G.Y. Lipopigment in the aging brain. Amer. Journal of Med. Genetics Supplement 5: 183-191, 1988.
416. Devine-Gage, E., Jenkins, E.C., Brown, T., Robakis, N.K., Wisniewski, H.M. The genetics of familial Alzheimer's disease. Age 11: 98-102, 1988.
417. Wisniewski, H.M., Currie, J.R., Brown, H.R., Barcikowska, M., Wegiel, J., Vorbrodt, A.W. Beta-peptide precursor protein producing and processing cells. In: The Molecular Biology of Alzheimer's Disease, edited by C.E. Finch and P. Davies. Cold Spring Harbor Lab., pp. 101-105, 1988.

418. Kascsak, R.J., Wisniewski, H.M. Pathogenesis of virus-induced and autoimmune nervous system injuries. In: Child Neurology and Developmental Disabilities, edited by J.H. French, S.H. Harel, and P.C. Casaer. Paul H. Brookes Publishing Co., pp. 89-98, 1988.
419. Currie, J.R., Wisniewski, H.M., Miller, D.L., Devine-Gage, E., Barcikowska, M., Wegiel, J., Mehta, P.D., Brown, H.R., and Vorbrodt, A. Pathogenesis and biochemistry of the Alzheimer amyloid beta-peptide. In: Senile Dementias, Current Problems in Senile Dementia, No. 2, II International Symposium, Rome, November 2-4, 1988, edited by A. Agnoli, J. Cahn, N. Lassen, and R. Mayeux, John Libbey Eurotext, Paris, pp. 15-25, 1988.
420. Brown, W.T., Rudelli, R.D., and Wisniewski, H.M. Fragile X syndrome: Neuropathology center. American J. of Med. Genetics 30: 201-205, 1988.
421. Robakis, N.K., Lahiri, D., Brown, H.R., Rubenstein, R., Mehta, P., Wisniewski, H.M., and Goller, N. Expression studies of the gene encoding the Alzheimer's disease and Down's syndrome amyloid peptide. In: Disorders of the Developing Nervous System: Changing Views on Their Origins, Diagnoses, and Treatments, edited by J.W. Swann, and M. Messer. Alan R. Liss, New York, pp. 183-193, 1988.
422. Rafalowska, J., Barcikowska, M., Wen, G.Y., and Wisniewski, H.M. Laminar distribution of neuritic plaques in normal aging, Alzheimer's disease and Down's syndrome. Acta Neuropath. 77: 21-25, 1988.
423. Kim, Y.S., Carp, R.I., Callahan, S.M., Wisniewski, H.M. Adrenal involvement in scrapie-induced obesity. Experimental Biology and Medicine 189: 21-27, 1988.
424. Wisniewski, H.M., and Iqbal, K. Paired helical filaments (PHF): Update 1988. In: Genetics and Alzheimer's Disease, edited by P.M. Sinet, Y. Lamour, and Y. Christen, Springer-Verlag, pp. 157-163, 1988.
425. Wisniewski, H.M., Merz, G.S., Rabe, A., Barcikowska, M., Moretz, R.C., and Devine-Gage, E.A. Current hypotheses of Alzheimer disease neuropathology and dementia. Drug Dev. Res. 15: 115-121, 1988.
426. Wen, G.Y., Rudelli, R.D., Kim, K.S., and Wisniewski, H.M. Tangles of ependyma-choroid plexus contain beta-amyloid protein epitopes and represent a new form of amyloid fiber. Archives of Neurology 45: 1298-1299, 1988.
427. Thormar, H., Brown, H.R., Goller, N.L., Barshatzky, M.R., and Wisniewski, H.M. Transmission of measles virus encephalitis to ferrets by intracardiac inoculation of a cell-associated SSPE virus strain. APMIS 96: 1125-1128, 1988.
428. Sturman, J.A., and Wisniewski, H.M. Aluminum. In: Metal Neurotoxicity, edited by Stephen C. Bondy and Kedar N. Prasad, CRC Press, Inc., Boca Raton, FL, pp. 61-85, 1988.
429. Rudelli, R.D., Wisniewski, H.M., and Barcikowska, M. Alzheimer neuropathology and clinical disease in non-Down syndrome mental retardation. In: Working Papers on Aging and Mental Retardation, Proceedings of the 111th Annual Meeting of the American Association on Mental Retardation, Los Angeles, CA, pp. 155-174, 1988.
430. Wisniewski, H.M. Recent advances and prospects in the brain sciences. In: Disability Unbound - The Challenge to the Brain and Information Sciences and Technologies, edited by D.M. Cone and D.P. Galamaga. Proceedings of a symposium held November 4, 1987 at the Rhode Island Medical Center in Cranston, Rhode Island, Published in Rhode Island, pp. 25-35, 1988.

431. Lossinsky, A.S., Song, M.J., and Wisniewski, H.M. High voltage electron microscopic studies of endothelial cell tubular structures in the mouse blood-brain barrier following brain trauma. Acta Neuropath. 22: 1-9, 1989.
432. Bancher, C., Lassmann, H., Budka, H., Jellinger, K., Grundke-Iqbal, I., Iqbal, K., Wiche, G., Seitelberger, F., Wisniewski, H.M. An antigenic profile of Lewy bodies: Immunocytochemical indication for protein phosphorylation and ubiquitination. J. of Neuropath. and Experimental Neurology 48: 81-93, 1989.
433. Wisniewski, H.M., and Sturman, J.A. Neurotoxicity of aluminum. In: Aluminum and Health. A Critical Review, edited by Hillel J. Gitelman, Marcel Dekker, Inc., New York and Basel, pp. 125-165, 1989.
434. Bancher, C., Brunner, C., Lassmann, H., Budka, H., Jellinger, K., Wiche, G., Seitelberger, F., Grundke-Iqbal, I., Iqbal, K., and Wisniewski, H.M. Accumulation of abnormally phosphorylated tau precedes the formation of neurofibrillary tangles in Alzheimer's disease. Brain Res. 477: 90-99, 1989.
435. Schupf, N., Silverman, W., Zigman, W.B., Moretz, R.C., and Wisniewski, H.M. Aluminum and Alzheimer's disease. The Lancet, p. 267, 1989.
436. Barcikowska, M., Silverman, W., Zigman, W., Kozlowski, P., Kujawa, M., Rudelli, R., and Wisniewski, H.M. Alzheimer-type neuropathology and clinical symptoms of dementia in mentally retarded people without Down syndrome. American Journal on Mental Retardation 93: 551-557, 1989.
437. Bancher, C., Grundke-Iqbal, I., Iqbal, K., Kim, K.S., and Wisniewski, H.M. Immunoreactivity of neuronal lipofuscin with monoclonal antibodies to the amyloid beta-protein. Neurobiology of Aging 10: 125-132, 1989.
438. Grundke-Iqbal, I., Iqbal, K., George, L., Tung, Y-C, Kim, K.S., and Wisniewski, H.M. Amyloid protein and neurofibrillary tangles coexist in the same neuron in Alzheimer disease. Proc. Natl. Acad. Sci. USA 86: 2853-2857, 1989.
439. Lossinsky, A.S., Song, M.J., Pluta, R., Moretz, R.C., and Wisniewski, H.M. Combined high-voltage and scanning electron microscopy on the same brain tissue samples for the study of blood-brain barrier injury. In: Proceedings of the 47th Annual Meeting of the Electron Microscopy Society of America, edited by G.W. Bailey, San Francisco Press, Inc., San Francisco, CA, pp. 986-987, 1989.
440. Wisniewski, H.M., Wen, G.Y., and Kim, K.S. Comparison of four staining methods on the detection of neuritic plaques. Acta Neuropath. 78: 22-27, 1989.
441. Chatterjee, N., and Wisniewski, H.M. A spirohydantoin derivative of oxymorphone: An agonist with delayed antagonist activity. Pharmacology Biochem. & Behavior 32: 939-943, 1989.
442. Korthals, J.K., Gieron, M.A., and Wisniewski, H.M. Nerve regeneration patterns after acute ischemic injury. Neurology 39: 932-937, 1989.
443. Barcikowska, M., Wisniewski, H.M., Bancher, C., and Grundke-Iqbal, I. About the presence of paired helical filaments in dystrophic neurites participating in the plaque formation. Acta Neuropath. 78: 225-231, 1989.
444. Wisniewski, H.M., Bancher, C., Barcikowska, M., Wen, G.Y., and Currie, J. Spectrum of morphological appearance of amyloid deposits in Alzheimer's disease. Acta Neuropath. 78: 337-347, 1989.

445. Brown, H.R., Goller, N.S., Rudelli, R.D., Dymecki, J., and Wisniewski, H.M. Postmortem detection of measles virus in non-neural tissue in subacute sclerosing panencephalitis. Annals of Neurology 26: 263-268, 1989.
446. Lossinsky, A.S., Badmajew, J.A., Robson, R.C., Moretz, R.C., and Wisniewski, H.M. Sites of egress of inflammatory cells and horseradish peroxidase transport across the blood-brain barrier in a murine model of chronic relapsing experimental allergic encephalomyelitis. Acta Neuropath. 78: 359-371, 1989.
447. Bancher, C., Grundke-Iqbal, I., Iqbal, K., Kim, K.S., and Wisniewski, H.M. A 31 kilodalton beta-protein immunoreactive polypeptide in neuronal lipofuscin. In: Alzheimer's Disease and Related Disorders, edited by K. Iqbal, H.M. Wisniewski, and B. Winblad. Alan R. Liss, Inc., New York pp. 913-924, 1989.
448. Wisniewski, H.M., Currie, J.R., Brown, H.R., Barcikowska, M., Wegiel, J., and Vorbrodt, A.W. Amyloidogenesis in Alzheimer disease and scrapie. In: Alzheimer's Disease and Related Disorders, edited by K. Iqbal, H.M. Wisniewski, and B. Winblad. Alan R. Liss, Inc., New York, pp. 869-876, 1989.
449. Bancher, C., Brunner, C., Lassmann, H., Budka, H., Jellinger, K., Seitelberger, F., Grundke-Iqbal, I., Iqbal, K., and Wisniewski, H.M. Tau and ubiquitin immunoreactivity at different stages of formation of Alzheimer neurofibrillary tangles. In: Alzheimer's Disease and Related Disorders, edited by K. Iqbal, H.M. Wisniewski, and B. Winblad. Alan R. Liss, Inc., New York, pp. 837-848, 1989.
450. Iqbal, K., Wang, G.P., Grundke-Iqbal, I., and Wisniewski, H.M. Laboratory diagnostic tests for Alzheimer's disease. In: Alzheimer's Disease and Related Disorders, edited by K. Iqbal, H.M. Wisniewski, and B. Winblad. Alan R. Liss Inc., New York, pp. 679-687, 1989.
451. Jenkins, E.C., Devine-Gage, E.A., Yao, X-L, Houck, G.E., Brown, W.T., Robakis, N.K., Wisniewski, K.E., Silverman, W.P., Reisberg, B., and Wisniewski, H.M. Beta-amyloid protein probe hybridized to chromosome 9 in 3 Alzheimer disease individuals. In: Alzheimer's Disease and Related Disorders, edited by K. Iqbal, H.M. Wisniewski, and B. Winblad. Alan R. Liss, Inc., New York, pp. 269-275, 1989.
452. Wisniewski, H.M. Milestones in the history of Alzheimer disease research. In: Alzheimer's Disease and Related Disorders, edited by K. Iqbal, H.M. Wisniewski, and B. Winblad. Alan R. Liss, Inc., New York, pp. 1-11, 1989.
453. Wisniewski, H.M., Iqbal, K., Bancher, C., Miller, S., and Currie, J. Cytoskeletal protein pathology and the formation of beta-amyloid fibers in Alzheimer's disease. Neurobiology of Aging 10: 409-412, 1989.
454. Madrid, R.E., and Wisniewski, H.M. Ultrastructural correlation with teased-fiber appearances in axonal degeneration and regeneration of guinea pig spinal roots. In: Peripheral Nerve Development and Regeneration, edited by E. Scarpini, M.G. Fiori, D. Pleasure, and G. Scarlato. Springer Verlag, Liviana Press, Padova, pp. 241-247, 1989.
455. Wisniewski, H.M., Rabe, A., Zigman, W., and Silverman, W. Editorial. Neuropathological Diagnosis of Alzheimer Disease. J. Neuropath. Exp. Neurol. 48: 606-609, 1989.
456. Carp, R.I., Kascsak, R.J., Wisniewski, H.M., Merz, P.A., Rubenstein, R., Bendheim, P., and Bolton, D. The nature of the unconventional slow infection agents remains a puzzle. Alzheimer Disease and Assoc. Disorders 3: 79-99, 1989.
457. Wisniewski, H.M., Wegiel, J., Wang, K.C., Kujawa, M., and Lach, B. Ultrastructural studies of the cells forming amyloid fibers in classical plaques. Can. J. Neurol. Sci. 16: 535-542, 1989.

458. Wisniewski, H.M., Mehta, P.D., Kim, K.S., and Merz, G.S. Cerebrospinal fluid-based laboratory test for Alzheimer's disease. In: Biological Markers of Alzheimer's Disease, edited by F. Boller, R. Katzman, A. Rascol, J.L. Signoret, and Y. Christen. Springer-Verlag Berlin, Heidelberg, pp. 23-29, 1989.
459. Schupf, N., Zigman, W.B., Silverman, W.P., Rabe, A., and Wisniewski, H.M. Genetic epidemiology of Alzheimer's disease. In: Aging Brain and Dementia: New Trends in Diagnosis and Therapy, edited by L. Battistin and F. Gerstenbrand, Alan R. Liss, Inc., Volume 54, pp. 57-58, 1990.
460. Vorbrodt, A.W., Dobrogowska, D.H., Lossinsky, A.S., Wisniewski, H.M. Changes in the distribution of anionic sites in brain micro-blood vessels with and without amyloid deposits in scrapie-infected mice. Acta Neuropath. 79: 355-363, 1990.
461. Kim, Y.S., Carp, R.I., Callahan, S.M., Natelli, M., and Wisniewski, H.M. Vacuolization, incubation period and survival time analyses in three mouse genotypes injected stereotactically in three brain regions with the 22L scrapie strain. J. of Neuropath. and Exp. Neurol. 49: 106-113, 1990.
462. Kim, Y.S., Carp, R.I., Callahan, S.M., and Wisniewski, H.M. Pathogenesis and pathology of scrapie after stereotactic injection of strain 22L in intact and bisected cerebella. J. of Neuropath. and Exp. Neurol. 49: 114-121, 1990.
463. Salerno, C., Wisniewski, H.M., and Rudelli, R.D. Effect of poppy seed ingestion on the TDx opiates assay. Therapeutic Drug Monitoring 12: 210-213, 1990.
464. Vorbrodt, A.W., Lossinsky, A.S., Dobrogowska, D.H., and Wisniewski, H.M. Sequential appearance of anionic domains in the developing blood-brain barrier. Devpmntl. Brain Res. 52: 31-37, 1990.
465. Rabe, A., Wisniewski, H.M., Schupf, N., and Wisniewski, H.M. Relationship of Down's syndrome to Alzheimer's disease. In: Application of Basic Neuroscience to Child Psychiatry, edited by S.I. Deutsch, A.W. Weizman, and R. Weizman. Plenum Publishing Corp., pp. 325-340, 1990.
466. Wisniewski, H.M., Moretz, R.C., Sturman, J.A., Wen, G.Y., and Shek, J.W. Aluminum neurotoxicity in mammals. Environ. Geochem. and Health 12: 115-130, 1990.
467. Moretz, R.C., Iqbal, K., and Wisniewski, H.M. Microanalysis of Alzheimer disease NFT and plaques. Environ. Geochem. and Health 12: 15-16, 1990.
468. Wisniewski, H.M., Rabe, A., Silverman, W., and Zigman, W. Reply to William I. Rosenblum's Letter to the Editor on Neuropathological Diagnosis of Alzheimer disease. J. Neuropathol. Exp. Neurol. 49: 189-190, 1990.
469. Brown, H.R., Goller, N.L., Rudelli, R.D., Merz, G.S., Wolfe, G.C., Wisniewski, H.M., and Robakis, N.K. The mRNA encoding the scrapie agent protein is present in a variety of non-neuronal cells. Acta Neuropath. 80: 1-6, 1990.
470. Iqbal, K., Grundke-Iqbal, I., and Wisniewski, H.M. Biochemical markers of Alzheimer's disease. In: Alzheimer's and Parkinson's Disease, edited by H.J. Altman and B.N. Altman. Plenum Press, New York, pp. 11-17, 1990.

ATTACHMENT 2

MANGANESE : COMMENTS ON HUMAN HEALTH RISK

Robert R. LAUWERYS, MD, MSc, DSc (Toxicology)

Professor of Industrial Toxicology and Occupational
Medicine

Director, Unit of Industrial Toxicology and
Occupational Health

University of Louvain

30.54. Clos Chapelle-aux-Champs

1200 Brussels

Belgium

For the general population, food usually constitutes the major source of manganese intake. The daily intake from food may vary over a wide range. In 1982, we have performed a duplicate meal study in Belgium and found values ranging from 0.6 to 8.8 mg manganese in 24-hr diets (median : 2.6). Drinking water generally contains less than 100 µg manganese/l (median value around 5 µg/l) but some mineralized water may contain higher concentrations. The median intake via drinking water is about 8 µg/day but can be as high as 2 mg/day from some water supplies. In many countries legislation requires that drinking water does not contain more than 50 µg manganese per liter. The adequate daily oral intake of manganese has been estimated at 2-3 mg but an additional amount of manganese (usually 1 mg po daily) is frequently prescribed during pregnancy in association with other oligoelements and various vitamins. Manganese absorption from the gastrointestinal tract is controlled by homeostatic mechanisms and varies from 1 to 5% (3% on the average). Individuals ingesting 3 mg manganese with food and drinking water will thus absorb on the average 90 µg manganese daily (range : 30 to 150 µg).

For non occupationally exposed subjects, manganese in ambient air represents a minor source of exposure. In rural and urban areas without significant manganese pollution, annual average levels of manganese in ambient air are mainly in the range of $0.005 - 0.07 \mu\text{g}/\text{m}^3$. With local industrial pollution, this level can rise to an annual average of $0.2 - 0.3 \mu\text{g}/\text{m}^3$ or even higher. Alveolar deposition of manganese may be estimated to be in the range of $0.07 - 0.5 \mu\text{g}/\text{day}$ as an average and $6-7 \mu\text{g}/\text{day}$ under high environmental exposure conditions. Even if the total amount deposited in the lungs is absorbed, the amount of manganese entering the circulation daily from the respiratory tract is much lower than that absorbed from the gastrointestinal tract.

Manganese is an essential element for mammals. It is incorporated in various enzymes and is a cofactor for a number of enzymatic reactions. Toxic effects of manganese have been mainly reported in workers chronically exposed to manganese dust and fume and more rarely among subjects living near ferromanganese plants (Elstad D. Norsk magazin for laegeridenskaben 3, 2527, 1939; Nogawa K. et al. Jpn J. Public Health 20, 315, 1973; Saric et al. International Conference on Heavy Metals in the Environment, vol. III, 27, 1975; Suzuki Y., Ind. Med. 12, 529, 1970) or drinking water contaminated by manganese (Kondakis et al. Arch. Environ. Health 44, 175, 1989). The two main target organs following chronic excessive exposure are the lungs (increased incidence of pneumonitis, bronchitis and chronic non specific lung disease) and the

central nervous system (neurobehavioral symptoms and neurological signs characteristic of an extrapyramidal syndrome).

The level of exposure to manganese which prevents the occurrence of central nervous system and lung disturbances has not yet been well defined. A WHO study group has concluded that there are indications that symptoms and signs which are not specific but may be connected with the early stage of the neurotoxic effects of manganese (manganism) can be found in a number of workers who have been exposed to manganese concentration of about 0.5 mg/m^3 of air; at approximately the same exposure levels, certain susceptible individuals may develop chronic manganese poisoning. Our recent studies on two cohorts of workers exposed to manganese oxides and salts support this conclusion. Adverse effects on the lungs of manganese exposed workers do not seem to appear at concentrations below $0.3\text{--}0.5 \text{ mg/m}^3$ of air. Studies performed on populations living near manganese emitting factories have suggested that an increased incidence of pneumonia and/or acute bronchitis may occur at lower levels of airborne manganese (references mentioned on page 2). It has been pointed out, however, that the increased incidence of pulmonary diseases found in these studies is not necessarily attributable to manganese itself. Other factors, including socioeconomic factors, which had not been considered may have influenced the results.

For occupational exposure, WHO (Geneva) has tentatively recommended a value of 0.3 mg of respirable manganese particles per m^3 of air as a

time-weighted average exposure. The preliminary results of an epidemiologic study among Belgian workers exposed to manganese dioxide in a dry alkaline-battery plant indicate that such exposure level may not necessarily prevents the occurrence of discrete neurofunctional changes in some subjects. This more recent study suggests that for occupational exposure it seems indicated to keep the airborne concentration (TWA) of respirable manganese particles below $150 \mu\text{g}/\text{m}^3$ (as manganese).

For the general environment, WHO (Air quality guidelines for Europe, WHO regional publications. European series; N° 23 - WHO Regional Office for Europe, Copenhagen 1987) has recommended an annual average of $1 \mu\text{g}$ manganese/ m^3 as a guideline value. This value should incorporate a sufficient margin of protection for the most sensitive population group.

March 1990

CURRICULUM VITAE

Robert R. LAUWERYS

Address

- Work : Department of Occupational Medicine and Hygiene
Université Catholique de Louvain - Medical School
30 Clos Chapelle-aux-Champs bte 30.54
1200 Brussels - Belgium
Tel : 02/764.32.20
- Home : 84 Nachtegalenlaan
1980 Tervueren - Belgium

Personal history

Born : 13.02.1938

Married

3 children

Educational history

- 1962 : M.D. University of Louvain - Belgium
- 1965 : Master of Industrial Health - University of Louvain - Belgium
- 1966 : Master of Science in Hygiene (Toxicology)
Harvard University, Boston, USA
- 1968 : Doctor of Science in Hygiene (Toxicology)
Harvard University, Boston, USA

Positions

- 1962 - 1965 : Clinical assistant - Internal Medicine Department
University Hospital - Louvain - Belgium

- 1966 - 1968 : Research fellow : Harvard University (School of Public Health), Boston, U.S.A. (CRB Graduate fellow and U.S. Public Health Service fellow)
- 1968 - 1969 : Research assistant : Toxicology Research Unit
Medical Research Council - Carshalton - U.K.
(Royal Society postdoctoral research fellow)
- Presently : - Professor of Occupational Health and Industrial Toxicology
Université Catholique de Louvain - Medical School - Brussels
Belgium
- Director Industrial Toxicology and Occupational Health Unit
School of Medicine - Catholic University of Louvain
 - Director Department of Occupational Medicine and Hygiene
Catholic University of Louvain

Scientific distinctions

- . Delta Omega (U.S. Public Health Honorary degree)
- . Prix Melsens (Académie Royale de Médecine de Belgique)
- . Member Royal Medical Academy of Belgium

Member

- National organizations
 - . Council on Hygiene - Ministry of Health, Belgium
 - . Scientific Committee - Work Compensation Fund - Ministry of Social Affairs, Belgium
 - . Société Belge de Médecine Interne
 - . Cercle des Alumni de la Fondation Universitaire

- International organizations

- . Council of Experts WHO
- . Advisory Subgroup on Toxicology of the European Medical Research Council (1974-1988)
- . CEC scientific committee on occupational exposure limits
- . Toxicology section of CEC - Scientific Advisory Committee to examine the toxicity and ecotoxicity of chemical compounds
- . Scientific Committee of "Centre d'Etudes et de Recherche des Charbonnages de France" from 1988
- . European Society of Toxicology
- . British Occupational Hygiene Society
- . Society of Toxicology, USA
- . American College of Occupational Medicine
- . American Industrial Hygiene Association
- . American Conference of Governmental Industrial Hygienists
- . International Commission on Occupational Health
- . Occupational Health in the Chemical Industry (Medichem)
- . Participation at many international committees involved in the assessment of health risk due to environmental and/or industrial pollutants and in the guidance of research in this field

Member of the Editorial Board of

- . Toxicology
- . International Archives of Occupational and Environmental Health
- . American Journal of Industrial Medicine (Foreign contributing editor)
- . Toxicology Letters
- . The Science of the Total Environment (from 1983 to 1987)
- . Archives des Maladies Professionnelles

- . Journal of Applied Toxicology
- . Journal de Toxicologie Clinique et Expérimentale
- . Archives of Environmental Health
- . Toxicology and Applied Pharmacology (from 1976 to 1982)
- . Cahiers des Notes Documentaires (INRS-France)
- . Biological Monitoring
- . Environmental Research
- . Environmental Management and Health

Activities

Teaching industrial toxicology

Research in industrial toxicology

- . study of the mechanism of action and the metabolism of industrial chemicals with the aim of developing methods for the early detection of adverse effects and assessing acceptable exposure levels
- . epidemiological and experimental approaches
- . main areas of research : metals, solvents, nephrotoxicity, biological monitoring of exposure.

Scientific publications :

Papers : 400 (English and French)

Books :

- * Toxicologie Industrielle et Intoxications Professionnelles, Masson, Paris, 1990 (3rd edition) - First edition translated in Italian and in Finnish
- * Industrial Chemical Exposure - Guidelines for Biological Monitoring. Biomedical Publications, Davis, California, 1983.

World Health Organization
Regional Office for Europe
Copenhagen



ATTACHMENT 3

MINISTÈRE DE LA SANTÉ PUBLIQUE
MINISTERIE VAN VOLKSGEZONDHEID
BULE - 1000
Rue de la Vierge Wytsmanstraat 14
1050 BRUXELLES

M. E.
J. M.
11017

Air quality guidelines for Europe

WHO Regional Publications, European Series No. 23

Manganese

General Description

Sources

Manganese (Mn) is an element widely distributed in the earth's crust. It is considered to be the twelfth most abundant element and the fifth most abundant metal. Manganese does not occur naturally in its native state. Oxides, carbonates and silicates are the most important among manganese-containing minerals. The most common manganese mineral is pyrolusite (MnO_2), usually mined in sedimentary deposits by open-cast techniques. Manganese occurs in most iron ores. Its content in coal is in the range 6-100 $\mu\text{g/g}$; it is also present in crude oil, but at substantially lower concentrations (1).

Manganese is mainly used in metallurgical processes, as a deoxidizing and desulfurizing additive and as an alloying constituent. It has other uses, e.g. in the production of dry-cell batteries, the production of manganese chemicals and in some other chemical processes, as well as in the manufacture of glass, in the leather and textile industries, and as a fertilizer. Organic carbonyl compounds are used as fuel-oil additives, smoke inhibitors, and as anti-knock additives in petrol (2).

Crustal manganese enters the atmosphere by a number of natural and anthropogenic processes, which include the suspension of road dusts by vehicles, wind erosion and suspension of soils, particularly in agricultural and building activities and quarrying processes. The resulting mechanically generated aerosols consist primarily of coarse particles ($\geq 2.5 \mu\text{m}$ mass median aerodynamic diameter (MMAD)). The smelting of natural ores and the combustion of fossil fuels also result in the ejection of crustal manganese to the atmosphere in the form of fume or ash in the fine-particle range ($\leq 2.5 \mu\text{m}$ MMAD). Manganese is released to the atmosphere during the manufacture of ferroalloys and other industrial processes. Nearly one half of all industrial and combusive emissions of manganese are from ferroalloy manufacture and about one tenth from fossil-fuel combustion. Minor amounts are generated by other processes. The use of manganese fuel additives constitutes an additional source (2).

As an element of low volatility, manganese tends to settle out near sources of pollution, but fine particles containing manganese can be distributed very widely. The most common forms of manganese compounds in the coarse particulates of crustal origin are oxides or hydroxides of oxidation state +2, +3, +4, and manganese carbonate. The manganese emitted by metallurgical processes consists of oxides. The manganese from combusted methylcyclopentadienyl manganese tricarbonyl (MMT), used in some countries as a fuel additive, is emitted primarily as Mn_3O_4 . Minute amounts of organic manganese compounds may be present in ambient air under certain conditions. However, MMT is rapidly photodegraded to inorganic manganese in sunlight. The estimated half-time is 10–15 seconds (2).

Manganese dioxide can react with sulfur dioxide or nitrogen dioxide to form manganous sulfate and dithionate, or manganese nitrate, respectively. It has been shown that aerosols of manganous sulfate can catalyse the oxidation of atmospheric sulfur dioxide to sulfur trioxide, thus promoting the formation of sulfuric acid (2).

Occurrence in air

The natural level of manganese in air is low. Background concentrations of 0.05 to 5.4 ng/m^3 over the Atlantic Ocean (3), and 0.01 ng/m^3 at the South Pole (4) have been reported. A concentration of 0.006 $\mu g/m^3$ in air at a height of 2500 m and an annual average concentration of 0.027 $\mu g/m^3$ at 823 m were reported (5). The national air surveillance network of urban areas in the USA indicated an annual average manganese concentration of 0.033 $\mu g/m^3$ in 1982 (2). In two cities of the Federal Republic of Germany (Frankfurt am Main and Munich), annual mean concentrations of manganese ranged between 0.03 and 0.16 $\mu g/m^3$ (6), and in Belgium over the period 1972–1977 concentrations of manganese, expressed also as annual means, between 0.042 and 0.456 $\mu g/m^3$ were reported (7). The Environmental Agency Japan (8) reported an annual mean manganese concentration in the air of Japanese cities of about 0.02–0.80 $\mu g/m^3$, with maximum 24-hour concentrations of 2–3 $\mu g/m^3$.

From these and other data it can be concluded that annual average levels of manganese in ambient air in nonpolluted areas range from approximately 0.01 to 0.03 $\mu g/m^3$, while in urban and rural areas without significant manganese pollution annual averages are mainly in the range of 0.01–0.07 $\mu g/m^3$. With local pollution near foundries, this level can rise to an annual average of 0.2–0.3 $\mu g/m^3$ and, in the presence of ferro- and silico-manganese industries, to over 0.5 $\mu g/m^3$ (9). In such places the average 24-hour concentrations may exceed 10 $\mu g/m^3$.

About 80% of manganese emitted into the atmosphere is associated with particles with an MMAD of less than 5 μm , and about 50% with particles with an MMAD of less than 2 μm . More recent data, however, tend to indicate that less than 50% of the total measured manganese in ambient air is found in fine particles (9).

Atmospheric particulate matter, including manganese, is transported by air currents until it is lost from the atmosphere by either dry or wet deposition. Manganese deposit in dustfall is more than twice that in rainfall (2).

The highest values of manganese concentrations in the working environment have been reported from manganese mines, ore-processing plants, dry-cell battery plants, and ferromanganese plants. In mining operations, manganese concentrations up to 250 mg/m^3 or even higher have sometimes been found. In dry-cell battery plants and ferromanganese plants the concentrations of manganese in air are lower. Values up to $5-8 \text{ mg/m}^3$, but occasionally also higher — up to 20 mg/m^3 or even more — have been reported (10). An important point is that in ferromanganese plants, but also in dry-cell battery plants, the size distribution of manganese aerosols is such that small particles prevail absolutely, compared with mining operations, where a smaller proportion of respirable particles is usually encountered. There is also some evidence that aerosols formed by condensation may be more harmful than those formed by disintegration. Whether or not this is caused by differences in the distribution of particle size remains to be clarified (10).

Routes of Exposure

Air

Because of the low solubility of manganese oxides, only inhaled manganese particles small enough to reach the alveoli ($\leq 2.5 \mu\text{m}$ MMAD) are likely to enter the bloodstream. Alveolar deposition of manganese may be estimated to be in the range of $0.07-0.5 \mu\text{g/day}$ as an average and $6-7 \mu\text{g/day}$ under high-exposure conditions (2).

Drinking-water

Concentrations of manganese in fresh water may vary from less than one to several thousand $\mu\text{g/litre}$ (11). Drinking-water generally contains less than $100 \mu\text{g}$ manganese per litre. In 100 of the largest cities in the USA, 97% of the surveyed public water supplies contained concentrations below $100 \mu\text{g/litre}$.

Food

Manganese concentration in foodstuffs varies markedly, but on the whole food constitutes a major source of manganese intake for humans. The highest concentrations are found in some foods of plant origin, especially wheat and rice, with concentrations between 10 and 100 mg/kg . Polished rice and wheat flour contain less manganese, since most of it is in the bran. High concentrations of manganese have been found in tea leaves. Eggs, milk, fruits and meat generally contain less than 1 mg manganese per kg (2). In a study performed in Canada (12) it was estimated that, of people's total manganese intake via food, 54% came from cereals and 14% from potatoes, whereas meat, fish and poultry provided only 2% of manganese intake. However, there is a difference in manganese concentrations for the same items in different countries and areas. Thus, daily manganese intake by adults from food may vary over a rather wide range ($2-12 \text{ mg}$). The daily intake for children aged 3-5 years averages 1.4 mg/day and for children aged 9-13 years 2.18 mg/day (13). The daily intake of manganese by bottled and breastfed infants is very low because of the low concentrations of manganese in both breast-milk and cow's milk (14).

MANGANESE

265

Relative significance of different routes of exposure

Total human exposure to manganese may be estimated from information on the levels in air, water and diet. The degree of uptake of manganese by inhalation is dependent upon particle size because of the low solubility of manganese oxides. Deposition of manganese in the alveoli can be calculated from the ambient concentration and the fraction present in fine particles. Alveolar deposition of manganese at current ambient levels may be estimated at $0.07 \mu\text{g}/\text{day}$ as an average and $6-7 \mu\text{g}/\text{day}$ under high-exposure conditions (2). Thus, daily intake through inhalation generally constitutes less than 0.1% of the total daily intake by the general population and rarely exceeds 1%, even in heavily polluted areas (1). Estimates of total thoracic deposition, including particles deposited in the tracheobronchial region, range from about $0.26 \mu\text{g}/\text{day}$ (average) to $10 \mu\text{g}/\text{day}$ (high) (2).

Ingested manganese in diet is estimated to be $0.002-0.004 \text{ mg}/\text{kg}$ body weight per day in infants and $0.06-0.08 \text{ mg}/\text{kg}$ body weight per day in adults (1). The daily intake of manganese from food by adults appears to be 2-9 mg. In Europe and the USA studies suggest a likely range of 2-5 mg, while in countries where grain and rice make up a major portion of the diet, the intake is more likely to be in the range of 5-9 mg. The consumption of tea may substantially add to the daily intake (1). The median intake via drinking-water is about $0.008 \text{ mg}/\text{day}$, but can be as high as $2.0 \text{ mg}/\text{day}$ for some water supplies. However, the average daily intake of manganese with water is unlikely to constitute more than 1-2% of the total intake of manganese. The ingestion of particles cleared from the respiratory tract probably constitutes no more than $0.01 \text{ mg}/\text{day}$ under the highest ambient exposure conditions (2).

Kinetics and Metabolism

There are no quantitative animal data on absorption rates for inhaled manganese compounds. Mena et al. (15) found that, in 17 humans exposed to a nebulized solution of manganese chloride and in 4 humans exposed to manganese oxide in a similar fashion, 40-70% of the deposited amount was recovered in the faeces. Both compounds were labelled with ^{54}Mn .

Manganese absorption is controlled by homeostatic mechanisms. The absorption rate will depend on the amount ingested and on tissue levels of manganese. In mice and rats, absorption of ingested manganese varies between 1 and 3.5%. It seems that manganese is absorbed equally well throughout the small intestine.

An increase in the iron content of milk decreases the whole-body retention of orally administered ^{54}Mn in rats by a factor of 10 (16). Manganese interactions with other elements (cadmium, nickel, indium, rhodium, selenium) and ethanol at the level of gastrointestinal absorption were also observed.

Mena et al. (15) found that 11 normal individuals absorbed an average of 3% of a dose of $200 \mu\text{g}$ manganese chloride labelled with ^{54}Mn . Another human study (17) indicates that manganese absorption takes place by diffusion in iron overload states and by active transportation in the duodenum and jejunum in iron deficiency states.

Absorbed manganese is rapidly eliminated from the blood and at first concentrates mainly in the liver. Excess metal may be distributed to other tissues. Concentrations of manganese are characteristic of the individual tissues, and almost independent of the species. Manganese preferentially accumulates in tissues rich in mitochondria. It penetrates the placental barrier in all species and is secreted in milk (2).

The highest concentrations of manganese in man have been found in the liver, kidneys, endocrine glands, and in the small and large intestines. The total body-burden for manganese has been estimated at 8-10 mg; about 33% of this amount is found in muscle tissue and 20% in the liver (18).

In blood, manganese is bound to proteins. In the trivalent state it can bind to transferrin and in the divalent state to an α -macroglobulin. The organic compound MMT is rapidly metabolized. The distribution in general is similar to that seen after exposure to inorganic manganese (1).

The apparent absence of cases of chronic oral manganese toxicity could be attributed to the extremely efficient homeostatic mechanism, well documented in animal and human studies, which prevents the accumulation of manganese in the body and maintains systemic plasma manganese concentrations at constant levels (2, 19). Manganese is excreted primarily via biliary clearance (2). However, it is possible that the capacity of the liver to excrete higher amounts of inhaled or injected manganese is limited, resulting in an increase in the tissue retention of manganese and, consequently, in the manifestation of toxic effects (19, 20). In excessive exposure other gastrointestinal routes may participate. Urinary excretion is small; in humans it is less than 1 μ g/day. This means that only about 0.01% of the body-burden is excreted daily via that route. Manganese is also excreted via sweat, hair, placenta and milk.

The biological half-time of manganese depends on the body-burden of manganese. Experiments have shown that manganese elimination from the brain is slower than that from the whole body. In humans without occupational exposure to manganese the half-time is around 35 days (21).

Health Effects

Manganese is an essential element. It is a constituent of several enzymes and can also activate many enzymes. Manganese deficiency was described only once in a man given a synthetic diet in which manganese had been omitted by mistake. Among the symptoms and signs were dermatitis, pigment changes of hair, retarded hair growth and hypocholesterolaemia (1). Excessive exposure to manganese has been shown to cause toxic effects in animals and humans. As manganese is regarded as a metal with a relatively low toxicity, acute poisoning by manganese in humans is very rare (2).

Effects on experimental animals

A large number of studies on laboratory animals were performed in order to explain the mechanism underlying the neurotoxicity of manganese seen in exposed workers (2). Although animal experiments show that in chronic

MANGANESE

267

manganese intoxication the central dopaminergic system is disturbed, the mechanism of manganese neurotoxicity has not been elucidated (2,22).

Recent studies indicate that age may play a role in the extrapyramidal disturbances seen in human manganism, suggesting that the aging brain shows differing susceptibility vis-à-vis the toxic effects of manganese (2,23). However, some new studies raise the intriguing possibility that lifetime treatment with manganese has some beneficial effects on the aging brain (23). An accurate dose-response relationship for inhalation exposure and neurotoxicity is unobtainable at present from the available animal studies (2).

Inhalation studies of the pulmonary effects show the occurrence of acute respiratory effects (oedema and leukocyte infiltration) when the level of exposure exceeds 20 mg MnO_2 per m^3 (2). Mice and monkeys exposed to MnO_2 by inhalation showed pathological effects after chronic (10 months) exposure to 0.7 mg/ m^3 (24). Studies in which animals were exposed for a long period of time (66 and 40 weeks) to about 0.1 mg MnO per m^3 as Mn_2O_3 particles or aerosols of respirable particle size showed no respiratory effects, but the studies had several methodological deficiencies which reduce confidence in the negative results (25,26).

It is plausible that exposure to manganese may increase susceptibility to pulmonary infections by disturbing the normal mechanism of lung clearance. However, it seems that a primary inflammatory reaction can occur in the lung after exposure to MnO_2 , if the concentration is high enough, without the presence of pathogenic bacteria (27). Studies on the influence of manganese on susceptibility to bacterial infections showed increased morbidity and mortality rates in animals infected before, during or after exposure to manganese dioxide (28,29).

Effects on humans

The neurological disorder known as manganism may result from occupational exposure to manganese dusts and fumes. Symptoms and signs of manganism have often been compared with Parkinson's disease, but certain differences should be noted. Patients with Parkinson's disease show pronounced disturbances of motor behaviour, which include tremor observed at rest rather than during intentional motor activity as in manganism. Fully developed manganism causes severe rigidity, with the extremities showing the "cog-wheel" phenomenon (2,30). Manganism usually appears after a prolonged exposure of two or more years, but it may result from exposure to high concentrations of manganese for only a few months. The full clinical picture of chronic manganese poisoning has been reported mainly in manganese miners, but also in other occupations where exposure to manganese is high (2,30). The disease has been found less frequently at exposure levels below 5 mg manganese per m^3 . However, there are reports of signs and symptoms which may be connected with subclinical or early clinical stages of chronic manganese poisoning in workers exposed to 0.3-5 mg manganese per m^3 . In connection with the toxic effects of manganese, a marked individual susceptibility has been observed (10,30). The data available for identifying effects below 1 mg manganese particles per m^3 are equivocal

and inadequate. Furthermore, no good biological indicators of manganese exposure are available at present.

The toxic effects of manganese on the pulmonary system vary in type and severity. There are reports of humans developing pneumonia in occupational but also in ambient exposure to manganese. Increased incidence of pneumonia was observed in workers, mainly miners, exposed to manganese at levels higher than 5 mg/m^3 (2). In a more recent study (31), an increased incidence of pneumonia and bronchitis was found in workers exposed to manganese concentrations of $0.4\text{--}16 \text{ mg/m}^3$ in a factory producing manganese alloys.

Elsstad (32) noted an 8-fold increase in mortality from pneumonia and a 4-fold increase in pneumonia morbidity in the general population living near a ferromanganese plant in Norway. The air concentrations of manganese were measured only once and were reported to be $46 \mu\text{g/m}^3$ at a distance of 3 km from the plant. It was also reported that the incidence of pneumonia followed the rate of production of manganese alloys. Analyses of lung tissues from 11 persons who died from pneumonia showed manganese concentrations of $0.35\text{--}1.63 \mu\text{g/kg}$ wet weight. A higher prevalence of nose and throat symptoms, and lower values of lung function tests compared with controls, were observed in children attending a school situated near a ferromanganese plant (33), where the average manganese exposure was about $7 \mu\text{g/m}^3$ (range $3\text{--}11 \mu\text{g/m}^3$). The study involved several hundred children, had a participation rate of over 97% and documented monitored levels of settled manganese dust for several years. Effects of still lower levels of airborne manganese (about $1 \mu\text{g/m}^3$) have been claimed to occur in a population living near a manganese alloy plant, with an increased incidence of acute bronchitis over an observation period of 4 years. The incidence of pneumonia did not seem to exceed the expected values (34). Although sulfur dioxide concentrations were measured (annual means: $10\text{--}30 \mu\text{g/m}^3$), other factors, including the socioeconomic factor, which had not been considered, may have influenced the results. Chronic bronchitis has been reported to be more prevalent in workers exposed to manganese if they are smokers (35).

A number of effects of manganese in other organs and systems have been claimed on the basis of results obtained from animal experiments and from epidemiological and clinical studies. These include a decrease in systolic blood pressure values, an increased rate of spontaneous abortions, changes in erythropoiesis and granulocyte formation, disturbed excretion of 17-ketosteroids, and changes in the activity of some enzymes (2). Reports of impotence in a number of patients with chronic manganese poisoning are also common. However, these effects have not been observed consistently. In some cases the implications of the results of animal experiments for human health are uncertain and therefore they cannot be regarded as relevant in the assessment of the potential health risk of ambient exposure to manganese.

Evaluation of Human Health Risks

Exposure

In urban and rural areas without significant manganese pollution, annual averages are mainly in the range of $0.01\text{--}0.07 \mu\text{g/m}^3$; near foundries the level

can rise to an annual average of $0.2-0.3\mu\text{g}/\text{m}^3$ and, where ferro- and silicomanganese industries are present, to more than $0.5\mu\text{g}/\text{m}^3$ with individual 24-hour concentrations exceeding $10\mu\text{g}/\text{m}^3$.

Health risk evaluation

Manganese is both an essential and, at higher levels, a toxic element. In assessing the health impact of manganese exposure, the effect on the CNS and the lungs should be regarded as the most significant. The neurological disorder known as manganism has been reported in the context of occupational exposure to manganese, seldom at levels below $5\text{mg}/\text{m}^3$. Studies related to signs and symptoms which may be connected with a subclinical or early stage of manganese poisoning in workers exposed to concentrations below $1\text{mg}/\text{m}^3$ are equivocal or negative.

Respiratory symptoms seem to occur at lower levels of exposure to manganese, e.g. below $1\text{mg}/\text{m}^3$, than do neurological symptoms and signs. Therefore, respiratory effects may be considered to be critical in ambient exposure to manganese. Schoolchildren exposed to about $7\mu\text{g}$ manganese per m^3 (range $3-11\mu\text{g}/\text{m}^3$) emitted from a ferromanganese plant had an increased prevalence of respiratory symptoms. This level of $7\mu\text{g}/\text{m}^3$ may therefore be considered the lowest-observed-effect level (33). This conclusion is supported by a report of increased incidence of acute bronchitis at levels of about $1\mu\text{g}/\text{m}^3$ in a population living near a manganese alloy plant (34).

The available evidence indicates that the current manganese levels generally found in industrialized countries are not in the concentration range associated with potentially harmful effects.

Guidelines

Available data from epidemiological studies suggest that the lowest-observed-adverse-effect concentration of manganese is about $7\mu\text{g}/\text{m}^3$. It is assumed that below $1\mu\text{g}/\text{m}^3$ (annual average), adverse health effects of environmental exposure to manganese are not likely to occur and therefore an annual average of $1\mu\text{g}/\text{m}^3$ is recommended as a guideline value. This value incorporates a sufficient margin of protection for the most sensitive population group. As the critical effect is one of respiratory irritancy it is desirable to have a short-term guideline value, but the present data base does not allow such estimations.

References

1. *Manganese*. Geneva, World Health Organization, 1981 (Environmental Health Criteria, No. 17).
2. *Health assessment document for manganese*. Cincinnati, OH, US Environmental Protection Agency, 1984 (Report No. EPA-600/8-83-013F).
3. Duce, R.A. et al. Atmospheric trace metals at remote northern and southern hemisphere sites: pollution or natural? *Science*, 187: 59-61 (1975).

4. Zoller, W.H. et al. Atmospheric concentrations and sources of trace metals at the South Pole. *Science*, 183: 198-200 (1974).
5. Georgii, H.W. et al. Die Verteilung von Schwermetallen in reiner und verunreinigter Atmosphäre [Distribution of heavy metals in clean and polluted atmospheres]. *Staub, Reinhaltung der Luft*, 34: 15-17 (1974).
6. Georgii, H.W. & Müller, J. Schwermetallaerosole in der Großstadtluft [Heavy metal aerosols in the air of large cities]. *Schriftenreihe des Vereins für Wasser-, Boden- und Lufthygiene*, 42: 39-50 (1974).
7. Kretzschmar, J.G. et al. Heavy metal levels in Belgium: a five-year survey. *Science of the total environment*, 14: 85-97 (1980).
8. [Results of measurements of national air sampling stations during 1973 (fiscal year)]. Tokyo, Environmental Agency Japan, 1975 [in Japanese].
9. Pace, T.G. & Frank, N.H. Procedures for estimating probability of non-attainment of a PM₁₀ NAAQS using total suspended particulate or inhalable particulate data. Research Triangle Park, NC, US Environmental Protection Agency, 1983.
10. WHO Technical Report Series, No. 647, 1980 (WHO recommended health-based limits in occupational exposure to heavy metals).
11. WHO guidelines for drinking-water quality. Vol. 2. Health criteria and other supporting information. Geneva, World Health Organization, 1984, pp. 275-278.
12. Méranter, J.C. & Smith, D.C. The heavy metal content of a typical Canadian diet. *Canadian journal of public health*, 63: 53-57 (1972).
13. Schlöge, C. & Wortberg, B. Manganese in the diet of healthy preschool and schoolchildren. *Acta paediatrica scandinavica*, 61: 648-652 (1972).
14. McLeod, B.E. & Robinson, M.F. Dietary intake of manganese by New Zealand infants during the first six months of life. *British journal of nutrition*, 27: 229-232 (1972).
15. Mena, I. et al. Chronic manganese poisoning: clinical picture and manganese turnover. *Neurology*, 17: 128-136 (1967).
16. Kostal, K. et al. The effect of iron additive to milk on cadmium, mercury and manganese absorption in rats. *Environmental research*, 22: 40-45 (1980).
17. Thomson, A.B.R. et al. Interrelation of intestinal transport system for manganese and iron. *Journal of laboratory and clinical medicine*, 78: 642-655 (1971).
18. Sumino, K. et al. Heavy metals in normal Japanese tissues. Amounts of 15 heavy metals in 30 subjects. *Archives of environmental health*, 30: 487-494 (1973).
19. Kello, D. & Vrcic, H. Manganese toxicity and homeostasis. In: *Proceedings of the Second International Symposium on Trace Elements, Human Health and Hair Analysis*. Amsterdam, May 18-19, 1981. Amsterdam, Elsevier Scientific Publishers (in press).
20. Thompson, T.N. & Kloos, C.D. Presystemic elimination of manganese in rats. *Toxicology and applied pharmacology*, 64: 236-253 (1982).
21. Mena, I. et al. Chronic manganese poisoning: clearance of tissue manganese concentrations with persistence of the neurological picture. *Neurology*, 18: 376-382 (1968).

22. Seth, P.K. & Chandra, S.V. Neurotransmitters and neurotransmitter receptors in developing and adult rats during manganese poisoning. *Neurotoxicology*, 5: 67-76 (1984).
23. Lai, J.C.K. et al. Differences in the neurotoxic effects of manganese during development and aging: some observations on brain regional neurotransmitter and non-neurotransmitter metabolism in a developmental rat model of chronic manganese encephalopathy. *Neurotoxicology*, 5: 37-47 (1984).
24. Suzuki, Y. et al. Effects of the inhalation of manganese dioxide dust on monkey lungs. *Tokushima journal of experimental medicine*, 28: 119-125 (1978).
25. Coulston, F. & Griffin, T. *Inhalation toxicology of airborne particulate manganese in rhesus monkeys*. Washington, DC, US Environmental Protection Agency, 1977 (Report No. EPA-600/1-77-026).
26. Ulrich, C.E. et al. Evaluation of the chronic inhalation toxicity of a manganese oxide aerosol. III — Pulmonary function, electromyograms, limb tremor, and tissue manganese data. *American Industrial Hygiene Association journal*, 40: 349-353 (1979).
27. Bergström, R. Acute pulmonary toxicity of manganese dioxide. *Scandinavian journal of work, environment & health*, 3(Suppl. 1): 1-11 (1977).
28. Maigret, R.Z. et al. Potentiating effects of manganese dioxide on experimental respiratory infections. *Environmental research*, 11: 386-391 (1976).
29. Adkins, B., Jr. et al. Increased pulmonary susceptibility to streptococcal infection following inhalation of manganese oxide. *Environmental research*, 23: 110-120 (1980).
30. Barbeau, A. Manganese in extrapyramidal disorders. *Neurotoxicology*, 5: 13-35 (1984).
31. Sarić, M. *Biological effects of manganese*. Research Triangle Park, NC, US Environmental Protection Agency, 1978 (Report No. EPA-600/1-78-001).
32. Elstad, O. Manganholdig fabrikkrök som medvirkende årsak ved pneumoni-epidemier i en industribygd [Factory smoke containing manganese as a predisposing cause in pneumonia epidemics in an industrial district]. *Norsk magasin for lægevidenskapen*, 3: 2527-2533 (1939).
33. Nogawa, K. et al. [Epidemiological studies on disturbance of the respiratory system caused by manganese air pollution. Report 1: Effects on respiratory system of junior high school students]. *Japanese journal of public health*, 20: 313-326 (1973) (in Japanese).
34. Sarić, M. et al. Acute respiratory diseases in a manganese contaminated area. In: *Proceedings of the International Conference on Heavy Metals in the Environment*. Toronto, Institute for Environmental Studies, University of Toronto, 1975. pp. 389-398.
35. Sarić, M. & Lučić-Palašić, S. Possible synergism of exposure to airborne manganese and smoking habit in occurrence of respiratory symptoms. In: Walton, W.H., ed. *Inhaled particles IV*. Oxford & New York, Pergamon Press, 1977. pp. 773-779.

ATTACHMENT 4

W. CLARK COOPER, M.D.
 3687 MT. DIABLO BLVD. (SUITE 320)
 LAFAYETTE, CALIFORNIA 94549
 PHONE (415) 284-5850

July 11, 1990

Air Docket
 US Environmental Protection Agency
 Room M-1500
 401 M Street, SW
 Washington, DC 20460

COMMENTS ON THE WAIVER APPLICATION BY ETHYL CORPORATION FOR
 METHYLCYCLOPENTADIENYL MANGANESE TRICARBONATE (MNT) AS A GASO-
 LINE ADDITIVE.

These comments are submitted at the request of the Ethyl Corporation in support of its application for a waiver to permit the use of MNT as an additive to gasoline. I am an independent consultant based in Lafayette, California. In 1984 I wrote a comprehensive review of the scientific literature entitled "The Health Implications of Increased Manganese in the Environment Resulting from the Combustion of Fuel Additives. A Review of the Literature". It was published in the Journal of Toxicology and Environmental Health (Vol. 14, pp 23-46).

The key questions considered in this review were:

1. How would the increments of manganese intake predicted from MNT use relate quantitatively to normal background levels and to levels known to be toxic?
2. Are there differences in the absorption, distribution, and excretion of inhaled manganese, as contrasted with ingested manganese, which would make small increases in airborne manganese unusually hazardous?
3. Would individuals with iron-deficiency anemias be unusually susceptible because of increased absorption of Mn?
4. Are infants hypersusceptible, because of increased intestinal absorption and poorly developed blood-brain barriers to metals?
5. Are there effects, other than those on the central nervous system associated with high concentrations of Mn, that deserve consideration? These include acute respiratory disease, interference with hematopoiesis, reproductive problems, mutagenicity, and carcinogenicity.
6. From consideration of the above factors, is the use of MNT as a fuel additive acceptable in terms of the public's health? "

After reviewing all available environmental, experimental and epidemiologic evidence I reached the following conclusions:

1. The increments in manganese intake in humans resulting from the use of MNT would be within the physiologic range and far below those known to be toxic.
2. While there are differences in the absorption, distrib-

Cooper
7/11/90

ution, and excretion of manganese that is inhaled as contrasted with manganese that is ingested, the differences are relatively small. Animals that have inhaled manganese derived from the combustion of MMT in concentrations greatly in excess of any that would result from MMT's use as a gasoline additive have not shown toxic effects.

3. There is moderately increased absorption of Mn associated with iron deficiency anemias, but this related to ingested Mn. Any increments in ingested Mn from MMT use would be within the variations that normally occur from differing dietary intakes.

4. Very young experimental animals have increased intestinal absorption of Mn and poorly developed blood-brain barriers to metals. While this suggests that they might be hypersusceptible to central nervous system effects from manganese, the increments of Mn from MMT use would lie within a range to which they are already being exposed and far below concentrations where such hypersusceptibility would be operative.

5. There is also no evidence to support any discernible impact of minute increments of Mn from MMT on the respiratory tract, the cardiovascular system, hematopoiesis, or reproduction. Neither should mutagenic or carcinogenic effects be anticipated, in view of the fact that total Mn intakes would remain in the physiologic range essential to health.

6. In spite of the fact that there are gaps in our knowledge of the metabolism of manganese and its functions and effects in biologic systems, these are more than balanced by experimental studies with high concentrations of Mn derived from the combustion of MMT. The minute increments of Mn that would result from the use of MMT as a gasoline additive should not have any impact on the public's health "

As of July 1990 I am not aware of any new evidence to alter these conclusions; if anything they have been strengthened. A review by Abbott of the Australian Department of Health(1987) and another by the Health Effects Institute (1983) have been in essential agreement.

Some new studies made available in the course of the recent hearing were reassuring. Roels et al (1987a, 1987b) reported epidemiologic surveys of Belgian workers exposed to inorganic Mn. Using very sensitive tests for neurotoxicity and respiratory effects, they reported only marginal changes at time-weighted averages of about 1,000 $\mu\text{g}/\text{m}^3$ over long periods of time.

The contributions by the Environmental Defense Fund(Dr. Ellen Silbergeld), Dr. Herbert L. Needleman, and Dr. David Hall added no evidence to support their positions against the waiver. The first two were largely devoted to criticism of Ethyl Corporation obviously related to the witnesses' long-standing concerns about lead.

The analogies between lead and manganese drawn by Drs. Silbergeld and Needleman are not scientifically valid or relevant to the current issue. They disregard the fact that manganese is essential to human health (acknowledged by Dr. Silbergeld on page 19), while lead is not. They ignore the fact that the range between lead exposures causing unquestionable toxicity and those where effects are minuscule is of the

Cooper
7/11/90

order of 10:1 or less. On the other hand, for manganese conclusive neurotoxicity in occupationally-exposed workers are associated with exposures thousands of times greater than the concentrations expected to result from the use of MT as a gasoline additive. The latter are within the range of background intake.

Some specific points raised by the witnesses deserve brief discussion. Dr. Silbergeld on page 2 criticized omission from Ethyl Corporation's presentation of a report from the California Air Resources Board (1989), saying that it "demonstrates that increases in airborne and deposited manganese are related to additions of manganese to gasoline already permitted in California". She did not point out that the report provided very weak evidence, limited to a few sites in Southern California to support the above. The authors concluded that "vehicular emissions of manganese may account for a significant part of the total at urban sites in Southern California". Noteworthy in the report was the fact that nearly all the measured Mn levels were extremely low, the mean concentrations reported for the Central Valley, the San Francisco Bay Area, and Southern California being 30, 15, and 30 nanograms per cubic meter respectively. In nearly all sites the fraction attributed to vehicular emissions was much less than that from dust arising from the earth's crust.

Dr. Needleman's dramatic calculation of the total amount of manganese that would be contributed to the nation's air from the sale of 100 billion gallons of gasoline containing 31.2 mg of MT per gallon was not particularly alarming. The figure he arrived at was 30,000 kg. When much of the soil of the United States contains 1000 parts per million of manganese, 33 additional tons does not sound like very much.

The NIEHS contribution summarized toxicologic studies of MNT, the danger of which is unquestioned and which must be tightly controlled. However this toxicity, prior to addition to gasoline, is not at issue. Dr. Hall's comments were clearly of interest but in several areas did not consider evidence that is already in the record. His statement that epidemiologic studies on Mn have been primarily on humans exposed to large amounts of Mn-bearing dust/fumes (as in miners, metal workers, or grinders) is of course true. As he knows these have shown severe effects in workers exposed to 5,000 $\mu\text{g}/\text{m}^3$ of Mn or more. In the range 1000 $\mu\text{g}/\text{m}^3$ and up effects have been minor or equivocal. There are of course no studies in exposure ranges of 100 or 10 or 1 $\mu\text{g}/\text{m}^3$. How would anyone design an epidemiologic study in exposure levels comparable to our normal background?

A fundamental flaw in all of the arguments raised against the waiver has been a disregard for the importance of dose. Nearly anything is toxic in high enough concentrations. No one disputes the fact that premature infants have been made totally blinded by postnatal exposure to 100% oxygen. Children can be killed by sodium chloride in amounts that seem surprisingly small. Neurotoxic effects from extremely high doses of Mn should not lead to the erroneous assumption that doses within the range normal background levels are hazardous. Even if some was in the tetroxide form and this were definitely more hazardous, we would still be dealing with extraordinarily low doses.

Cooper
7/11/90

It is unfortunate that the opponents of the waiver have have paid so little attention to probable environmental benefits from the additive. The evidence for these seems far more convincing than that for hypothetical and improbable health risks from Mn.

I sincerely hope that the EPA considers the scientific evidence carefully in reaching its decision on the waiver.

Respectfully,



W. Clark Cooper, MD

PARTIAL LIST OF REFERENCES

Cooper, WC (1984) The health implications of increased manganese in the environment resulting from the combustion of fuel additives: a review of the literature. J. Toxicol. Environ. Health 14:23-46

Abbott, RJ (1987) Methylcyclopentadienyl manganese tricarbonyl (MMT) in petrol: The toxicological issues. The Science of the Total Environment 67: 247-255.

Davis, DW et al (1980) Origins of manganese in air particulates in California. JAPCA 32:1152-

Roels et al (1987a) Epidemiological survey among workers exposed to manganese: effects on lungs, central nervous system and some biological indices. Am.J. Ind. Med. 11:307-327.

Roels et al (1987b) Relationship between external and internal parameters of exposure to manganese in workers from a manganese oxide and salt producing plant. Am.J. Ind. Med 11:297-305.

The Health Effects Institute (1988: Potential health effects of manganese in emissions from trap-equipppd diesel vehicles. Report dated September 1988

U.S. Environmental Protection Agency (1990); Transcript of Hearing on Ethyl Corporation Fuel Waiver Application, June 22, 1990.

January 10, 1985

CURRICULUM VITAEName: W. Clark CooperDate of birth: June 22, 1912Place of birth: Manila, Philippine IslandsEducation: M.D., University of Virginia, 1934

M.P.H., Harvard School of Public Health, 1958

Positions held:

1934-1937	Interne and assistant resident in internal medicine, University Hospitals of Cleveland
1937	Resident fellow in nutrition, University of Cincinnati
1937-1940	First Lieutenant, U.S. Army Medical Corps
April 1940	Entered U.S. Public Health Service as Commissioned Officer
1941-1950	Research in chemotherapy of malaria, National Institute of Health, Bethesda, Maryland
1950-1952	Assistant Chief of Medicine, U.S. Public Health Service Hospital, San Francisco, California
1952-1957	Chief, Occupational Health Field Headquarters, Division of Occupational Health, PHS, Cincinnati, Ohio
1958-1961	Chief, Program Planning and Analysis, Division of Occupational Health, PHS, Washington, D.C. During part of this time also functioned as Acting Research Grants Administrator
Sept. 1961- April 1962	Deputy Chief, Division of Occupational Health and Chief, Program Planning and Analysis.
April 1962- August 1962	Acting Chief, Division of Occupational Health, Public Health Service, Washington, D.C.
August 1962- Sept. 1963	Chief, Division of Occupational Health, Public Health Service, Washington, D.C.
Sept. 1963	Retired as Medical Director, U.S. Public Health Service
Sept. 1963- Sept. 1972	Research Physician (Occupational Health) to Professor of Occupational Health in Residence, School of Public Health, University of California, Berkeley.
1972-1977	Vice President, Tabershaw/Cooper Associates, Inc., later Equitable Environmental Health, Inc.
1978 to present	Consultant in Occupational Health

Specialty Board Certification:

American Board of Internal Medicine, 1949
American Board of Preventive Medicine (Occupational Medicine), 1962
American Board of Industrial Hygiene (Comprehensive), 1962

Professional Organizations:

Fellow, American Association for the Advancement of Science
Fellow, American Public Health Association
Fellow, American College of Chest Physicians
(Member of Committee on Occupational Diseases of the Chest)
Fellow, American Occupational Medical Association
Fellow, American Academy of Industrial Hygiene
Fellow, Royal Society of Tropical Medicine and Hygiene (London) 1948-1980
Fellow, Royal Society of Medicine (London)
Member, American Society of Tropical Medicine and Hygiene 1942-1980
Member, Permanent Commission and International Association on
Occupational Health
Member, Western Occupational Medical Association
Member, American Academy of Occupational Medicine
Member, Alameda-Contra Costa Medical Association
Member, Society for Environmental and Occupational Health
Member, Society for Epidemiologic Research
Member, Cosmos Club, Washington, D.C.

Special Committees and Activities:

Member, Section on Environmental Health, American College of Chest Physicians
Member, Committee for Criteria Document Review, American Academy of Occupational Medicine
Member, Committee on Mineral Fibres, Permanent Commission and International Association on Occupational Health
Medical Consultant, U.S. Atomic Energy Commission, 1964-1973
Member, Panel on Clinical Testing of Antimalarials, Board of Coordination of Malarial Studies, 1944-1946
Member, Threshold Limits Committee of the American Conference of Governmental Industrial Hygienists, 1956-1961
Member, Committee on Biologic Effects of Atmospheric Pollutants, National Academy of Sciences, 1970-1972, and Chairman, Panel on Asbestos, 1970-1971
Member, Task Force on Chronic Bronchitis and Emphysema (USPHS and National Tuberculosis Association), 1966-1967
Medical Advisor, Western States Asbestos Health Plan (Western States Conference of Asbestos Workers and Associated Insulation Contractors of the Western States, 1968-1983
Associate Editor, Journal of Occupational Medicine, 1971-1976
Member, Committee on Public Information in the Prevention of Occupational Cancer, National Academy of Sciences, 1975-1977
Member, Subcommittee on Occupational Health, Committee on Military Environmental Research, National Research Council, 1979-1981

Selected publications and reports

Spies, TD & C Cooper: The diagnosis of pellagra. International Clinics 4:1-11, 1937

Spies, TC, C Cooper & MA Blankenhorn: The use of nicotinic acid in the treatment of pellagra. JAMA 110:622-27, February 26, 1938

(38 reports on chemotherapy of avian and human malaria, 1945-1953)

Coatney, GR, WC Cooper: Recrudescence and relapse in vivax malaria. Proceedings 4th International Congress Tropical Medicine and Malaria, Washington DC, 1948, pp 629-39

Cooper, WC: Summary of Antimalarial Drugs. Public Health Rep, 64: 717-32, June 10, 1949

Cooper, WC: Summary of Antimalarial Drugs. J Am Pharm Assoc, 10: 552-56, September 1949

Cooper, WC, LJ Cralley, WH Clark, BR Hubbard, DJ Hurley, RR Sullivan: Pneumoconiosis in Diatomite Mining and Processing. Public Health Service Publication No. 601, 1958, 96 pp

Cooper, WC: Case Study No. 1. Uranium Milling and Mining. Proceedings of the Conference of Epidemiologic Research in Occupational Health. J Occup Med (Special Supplement, pp 614-622), October 1962

Cooper, WC, PE Enterline, ET Worden: Estimating Occupational Disease Through Medical Care Plans. Public Health Rep, 77: 1065-70, December 1962

Cooper, WC, VB Perone, LD Scheel & RG Keenan: Occupational hazards from castor bean pomace: tests for toxicity. Am Ind Hyg Assn J, 25:431-438, 1964.

Tabershaw, IR, WC Cooper: Sequelae of Acute Organic Phosphate Poisoning. J Occup Med, 8:5-20, January 1966

Cooper, WC, IR Tabershaw: Biologic Effects of Nitrogen Dioxide in Relation to Air Quality Standards. Arch Environ Health, 12:522-30, April 1966

Apen, EM, WC Cooper, RJM Horton, LD Scheel: Health Aspects of Castor Bean Dust, Review and Bibliography. Public Health Service Publication No. 999-AP-36, 1967

Cooper, WC & IR Tabershaw: An appraisal of the potential effect on health of automatic pyrethrin dispensers. Prepared for C.A.P. Inc. May 1, 1967

Cooper, WC: Asbestos as a Hazard to Health, Fact and Speculation. Arch Environ Health, 15:285-90, September 1967

Cooper, WC: Uranium Mining and Lung Cancer. J Occup Med, 10:82-88, February 1968

Balzer, JL, WC Cooper: The Work Environment of Insulating Workers. Am Ind Hyg Assoc J, 29:222-27, May/June, 1968

Tabershaw, IR & WC Cooper: A Labor-Management Occupational Health Service in a Construction Industry. Arch Environ Health, 21:784-787, December 1970

Fowler, DP, JL Balzer, WC Cooper: Exposure of Insulation Workers to Airborne Fibrous Glass. Am Ind Hyg Assoc J, 32:86-91, February 1971

Balzer, JL, WC Cooper, DP Fowler: Fibrous Glass-Lined Air Transmission Systems, An Assessment of Their Environmental Effects. Am Ind Hyg Assoc J, 32:512-18, August 1971

Balzer, JL, DP Fowler, WC Cooper: Exposure of Sheet Metal Workers to Airborne Fibrous Glass; Report to Health and Safety Committee, National Insulation Manufacturers Association; University of California, Berkeley, School of Public Health, Division of Environmental Health Sciences, September 1971

Cooper, WC, IR Tabershaw, KW Nelson: Laboratory Studies of Workers in Lead Smelting and Refining. In Proceedings (of an) International Symposium (of the) Environmental Health Aspects of Lead, Amsterdam, October 2-6, 1972, Commission of the European Communities, Centre of Information and Documentation, Luxembourg, 1973, pp 517-530

Balzer, JL, DP Fowler, WC Cooper: Dust-Producing Potential of Construction Materials (1972). In Proceedings of a Symposium ... on Safety and Health in Shipbuilding and Ship Repairing, Helsinki, 1971, Geneva, International Labour Office, 1972, pp 107-22

Utidge, MD, WC Cooper: Sputum Cytology and Occupational Pulmonary Disease. J Occup Med, 15:253-54, March 1973

Cooper, WC: Indicators of Susceptibility to Industrial Chemicals. J Occup Med, 15:355-59, April 1973

Cooper, WC, WR Gaffey: Mortality of Workers in Lead Smelters and Lead Battery Plants. In Proceedings (of an) International Symposium on Recent Advances in the Assessment of the Health Effects of Environmental Pollution, Paris, June 24-28, 1974, Commission of the European Communities, Luxembourg, 1974

Cooper, WC and J Miedema: Asbestosis in the Manufacture of Insulating Materials, in Biological Effects of Asbestos, 1973, Lyon, International Agency for Research on Cancer, pp 175-178

Cooper, WC and W Gaffey: Mortality of Lead Workers. J Occup Med, 17: 100-107, February, 1975

Cooper, WC: Radiographic Survey of Perlite Workers. J Occup Med, 17:304-307, May 1975

Cooper, WC: Pulmonary Function in Perlite Workers. J Occup Med, 18:723-729, November, 1976

Cooper, WC: Cancer Mortality Patterns in the Lead Industry. Ann NY Acad Sci, 271:250-259, 1976

Cooper, WC and WR Gaffey: Mortality study of lead workers. Arhiv za Higijenu Rada I Toksikologiju (Zagreb), 26:209-229, 1976

Cooper, WC: Airborne asbestiform minerals in the non-occupational environment. Prepared for American Iron Ore Association, Inc., 1976

Cooper, WC: Airborne manganese and public health, with emphasis on the use of manganese-containing gasoline additives. Presented at Medichem Symposium, San Francisco, September 9, 1977

Cooper, WC: Epidemiologic evidence of the effect of type of asbestos and fiber dimension on the production of disease in man. Workshop on Asbestos: Definitions and Measurement Methods. U.S. Department of Commerce/National Bureau of Standards, NBS Publication 506, pp 121-132, November, 1978

Cooper, WC: Predictions of mortality from occupational cancer with special reference to asbestos. Presented at the American Industrial Hygiene Conference, Chicago, Illinois, May 30, 1979.

Cooper, WC, J Murchio, W Popendorf, HR Wenk: Chrysotile Asbestos in a California Recreational Area. Science 206:685-688, November 9, 1979

Cooper, WC, O Wong and L Kheifets: Mortality in employees of lead battery plants and lead production plants, 1947-1980. Prepared for International Lead Zinc Research Organization, August 1, 1980 (publication pending).

Cooper, WC: Health considerations in the perlite industry. Proceedings of the National Workshop on Substitutes for Asbestos, Arlington, Virginia, July 14-16, 1980, EPA 560/3-80-001 November, 1980, pages 505-510

Cooper, WC: Mortality in employees of lead production facilities and lead battery plants, 1971-1975. Environmental Lead, DR Lynam, LG Piantanida & JF Cole, Eds., pp 111-143, Academic Press, 1981.

Cooper, WC: Epidemiologic study of vinyl chloride workers: mortality through December 31, 1972. Environmental Health Perspectives 41:101-106, 1981.

Cooper, WC: The carcinogenic potential of ferric oxide, with particular reference to hematite: review of the evidence. Prepared for American Iron Ore Association, March 20, 1981.

Cooper, WC and O' Wong: A study of mortality in a population of nickel miners and smelter workers. Prepared for the Hanna Nickel Smelting Company, Riddle, Oregon, June 22, 1981.

Cooper, WC: Societal implications of occupational research. In Legal & Ethical Dilemmas in Occupational Health, JS Lee and WN Rom Eds., pp 269-276, Ann Arbor Science, 1982

Cooper, WC: Asbestos: protection in the workplace. Problems of medical surveillance. Presented at World Symposium on Asbestos, Montreal, Quebec, Canada, May 25, 1982

Cooper, WC: Mortality in employees of three plants which produced chromate pigments. Prepared for Dry Color Manufacturers Association, Inc., March 10, 1983.

Cooper, WC: The health implications of manganese in the environment resulting from the combustion of fuel additives. A review of the literature. J. Toxicology & Environmental Health, 14:23-46, 1984.

Cooper, WC & EN Sargent: A 26-year follow-up of workers in a diatomite mine and mill. JOM 26:456-460, 1984.

Cooper, WC: An epidemiologic study of workers in the taconite mining and milling industry. Prepared for the American Iron Ore Association, March 26, 1984.

Cooper, WC & EN Sargent: A study of chest radiographs and pulmonary function in perlite workers. Prepared for the Perlite Institute, Inc., May 1, 1984.

Cooper, WC: Mortality in employees of lead battery plants and lead production plants, 1947-1980. Final Report Project LH-157, prepared for International Lead Zinc Research Organization, Inc., August 1, 1984. Presented in summary form at XXI International Congress on Occupational Health, Dublin, Ireland, September 1984.

Cooper, WC, O Wong & L Kheifets: Mortality in employees of lead battery plants and lead producing plants, 1947-1980, submitted for publication, Scandinavian J of Work, Environment and Health, November 1984.

W. Clark Cooper, M.D.
3687 Mount Diablo Blvd.
Suite 320
Lafayette, CA 94549

(415) 284 5850

APPENDIX

8

Analytica Chimica Acta, 120 (1980) 171–176

© Elsevier Scientific Publishing Company, Amsterdam — Printed in The Netherlands

DETERMINATION OF METHYLCYCLOPENTADIENYLMANGANESE-TRICARBONYL BY GAS CHROMATOGRAPHY—ATOMIC ABSORPTION SPECTROMETRY AT ng m^{-3} LEVELS IN AIR SAMPLES

M. COE, R. CRUZ and J. C. VAN LOON

*Department of Geology, Chemistry and the Institute for Environmental Studies,
University of Toronto, Toronto, Ontario M5S 1A1 (Canada)*

(Received 2nd June 1980)

SUMMARY

A procedure is given for the determination of methylcyclopentadienylmanganese tricarbonyl (MMT) at ng m^{-3} concentrations in air. The method involves trapping of MMT in a small segment of gas chromatographic column and then determination by gas chromatography with an electrothermal atomic absorption detector. The detection limit of the procedure is 0.05 ng m^{-3} . Air samples from an underground car-park (when MMT was detected) were found to contain between 0.1 and 0.3 ng m^{-3} MMT. MMT was not detected in any of the street air samples taken.

Methylcyclopentadienylmanganesetricarbonyl (MMT) is added to many of the so-called "unleaded" gasolines sold in North America, as an antiknock compound. Levels of MMT in these gasolines average about 0.16 g gal^{-1} . MMT was discovered and patented by Ethyl Corporation and there is little information in the scientific literature concerning this compound. The current widespread distribution of MMT as a gasoline additive makes it essential that a method be developed for its determination in environmental samples. Most procedures for the determination of MMT are indirect relating it to the total manganese found in a sample. Turkel'traub et al. [1] described a gas chromatographic procedure for the related compound cyclopentadienylmanganesetricarbonyl (CMT). Only one procedure, that of Uden et al. [2], has been reported for the direct determination of MMT. This latter procedure is for the relatively high levels of MMT found in gasoline. In the following paragraphs, a procedure is outlined for the gas chromatographic/atomic absorption determination of MMT directly at ng m^{-3} levels in air.

The techniques of analytical atomic spectrometry can be used to advantage in the study of metal compounds, as metal specific detectors for chromatography. Uden et al. [2] used d.c. argon plasma emission spectrometry as the detector in their work. The detection limit obtainable by d.c. argon plasma emission for manganese is poorer than for electrothermal atomic absorption. Thus this latter detector was chosen in the current work because of the very low levels of MMT expected in environmental samples.

EXPERIMENTAL

Apparatus and reagents

A Perkin-Elmer 603 atomic absorption spectrometer was used with a Perkin-Elmer HGA 2100 furnace. A deuterium arc background corrector was employed. A Pye gas chromatograph (Series 104) was interfaced to the graphite furnace using a tantalum connector as previously described [3]. A glass chromatographic column (2.3 m long, 6 mm o.d.) was packed with 3% OV-1 on high-performance Chromosorb W (80/100 mesh). The gas from the chromatograph was transferred to the furnace through teflon-lined aluminium tubing.

The gas chromatographic set up is illustrated in Fig. 1. (A) connects to a nitrogen cylinder. (B) is the sample oven containing the sample trap and 4-way valve. (C) is the gas chromatograph connected through a standard injection valve to the column. (D) is the graphite furnace. The operating conditions for the gas chromatograph and the graphite furnace were as shown in Table 1.

Samples of air were collected in teflon-lined aluminium U-tubes (30 cm long, 3 mm o.d.) packed with 3% OV-1 on Chromosorb W (80/100 mesh). These tubes were placed in a water-ice cooling bath. Air entered through an air filter previously described [3] and was pumped through the U-tube trap at about 70 ml min^{-1} using a vacuum pump. The length of sampling time and the average flow rate, checked frequently during sampling, were used to compute sample volume.

MMT (Alfa Division, Ventron Corporation, Danvers, Massachusetts) was diluted in isooctane to produce the working standards. These standards were prepared fresh daily.

Procedures

Air samples were collected for various periods of up to 100 h and were retained in a freezer until run.

The gas chromatographic system is assembled as shown in Fig. 1. All transfer lines between the sample oven (B) and the gas chromatograph (C) and thence to the graphite furnace (D), must be wrapped in 36-W heating tape and held at 150°C throughout the procedure.




Fig. 1. Gas chromatograph system.

TABLE 1

Instrumental operating conditions

Gas chromatography		Atomic absorption spectrometry	
Carrier gas flow rate	80 ml min ⁻¹	Ash	300°C
Oven temperature program	115°C isothermal	Atomize	1800°C
Injection port temperature	150°C	Internal gas flow	0
Outlet and transfer tube temperature	150°C	External gas flow	60 ml min ⁻¹
		(N ₂)	
		Wavelength	279.5 nm
		Slit 4	(0.7 nm)
		Background correction mode,	
		Scale expansion (x 5)	

With the 4-way valve in the sample oven in the bypass position, the sample trap is placed in the cool sample oven and connected with Swagelok fittings. The nitrogen carrier gas flow rate is adjusted to 100 ml min⁻¹ (which reduces to 80 ml min⁻¹ when the sample trap is switched on). The sample oven is heated at 150°C for 10 min (no flow through the sample tube). The 4-way valve is then turned to start the flow through the sample tube, and the atomic absorption furnace program is initiated immediately. The gas chromatograph is run isothermally at 115°C. Gas flow is continued until the MMT peak has been recorded.

Calibration is done by injecting an appropriate μ l standard of MMT directly into the cooled, spent, sample tube in the sample oven. The above procedure is then followed exactly as for the sample.

RESULTS AND DISCUSSION

Optimization of the procedure

Collection of MMT. Because of the very low levels of MMT expected in air and other environmental samples, it was necessary to employ a method of preconcentration. Two possibilities exist: (1) the MMT can be trapped in a solvent contained in a bubbler; or (2) the MMT can be trapped on an adsorber held at very low temperature. Both of these strategies were tried.

A micro bubbler containing hexane was cooled in an ice bath. After collection of the sample the hexane volume was reduced to 0.4 ml. A 10- μ l aliquot was injected into the gas chromatograph. The detection limit using this approach was about 0.02 μ g m⁻³ of air for an 8-h sampling period. This was shown to be too poor for the present purposes.

The second trapping strategy is that outlined in the proposed procedure above. MMT is collected on a teflon-lined U-tube containing packing material similar to that used in the gas chromatographic columns. It was necessary to build a sample oven which contained a 4-way valve (which allows by-passing of the sample U-tube) so that any lines or other components carrying MMT

could be heated. This is because MMT has a relatively high boiling point and tends to condense out on cool transfer lines.

Decomposition of MMT in transfer lines. In a similar study of tetraalkyllead compounds, the problem of decomposition of these compounds in the metal transfer lines was noted. Teflon-lined tubing prevented this problem and was used as a precaution in this study as well.

Calibration. Ideally it would be desirable to inject standard solutions through the injection valve for calibration and avoid collection on the sample trap. However, the chromatographic peaks obtained by direct injection and those obtained using the sample trap had slightly different retention times and were differently shaped. Thus, it was necessary to inject the calibration standard onto the spent sample U-tube and follow the procedure exactly as for the samples. Peak area rather than peak height was used in the calculations.

Comparison of atomizers. Interfacing of gas chromatography and atomic absorption occurs through the atomizer. When best atomic absorption sensitivity is required, a commercial electrothermal atomizer (furnace or rod) is usually chosen. These devices are relatively expensive and are not available in all atomic absorption laboratories to be tied up for this purpose. Thus it was of interest to determine the relative detection limits obtainable for inexpensive homemade quartz-tube atomizers which were developed for a study of tetraalkyllead compounds [3].

These atomizers consist of a quartz tube wrapped with heating wire or a slotted quartz tube held in an air-acetylene flame. The relative detection limits of the wire-wrapped quartz tube and the Perkin-Elmer HGA 2100 were the same, whereas that of the slotted quartz tube was 20 times greater in comparative tests with MMT. Thus, if desired, a quartz tube wrapped in heating wire can be used in place of the commercial furnace with no penalty to be paid in detection limit.

It is interesting to note that MMT atomizes readily giving free manganese atoms at the relatively low temperature of 1000°C attainable with the heating wire wrapped quartz tube atomizer. This temperature is in marked contrast to 2700°C often recommended as the atomization temperature in the commercial graphite furnace atomizers. In this regard, it was found necessary, for maximum sensitivity, to align the manganese hollow-cathode lamp so that the optical beam just grazed the inner surface of the quartz tube nearest the gas inlet. Sensitivities obtained in the center of the tube are greatly reduced.

Analysis of air samples

Air samples of up to 15 m³ were taken at a variety of locations on the streets of Toronto. No MMT was detected in any sample (detection limit 0.05 ng m⁻³). This compares with 14 ng m⁻³ for total tetraalkyllead compounds found in street air in similar locations. Samples were then collected in an underground car-park beneath the Chemistry and Physics buildings at the University of Toronto. In a few of these samples MMT was detected.

Figure 2 shows the chromatograms obtained when a calibration standard




Fig. 2. (A) Chromatograph for standard sample containing MMT and CMT. (B) Chromatogram of an air sample taken in an underground car-park.

containing CMT which can be used as an internal reference if desired [2]) and MMT were run according to the proposed procedure. The level of MMT in the air sample was 0.3 ng m^{-3} . Other samples yielded values between 0.1 and 0.3 ng m^{-3} . It should be noted that the retention times of the MMT peak shown in the chromatogram for the air sample and the calibration standard are slightly different (ca. 0.2 min). It was not uncommon to have this variation which reflects the reproducibility problems found when sample traps are used.

It is interesting to speculate as to why the MMT levels in air are so much lower than those of tetraalkyllead when it is likely that a large fraction of the vehicles are using "unleaded" gasoline containing MMT. To this end, a cursory study of MMT in air was attempted.

MMT, liquid and vapor, was injected into clear 2-l glass bottles kept in the light and in the dark. Samples were then taken of the contained air and isooctane rinsings of the bottle walls at various intervals. Results of these studies suggest that MMT is quickly decomposed in air, more quickly in the light than in the dark. This conclusion must be treated as highly qualitative since a number of problems were encountered during this work.

Regarding this experiment, it was particularly difficult to keep the MMT suspended in the air. There is a great tendency for this compound to condense out on the sides of the bottles. Apart from making sampling difficult, this calls into question whether such a system is valid for determining the stability of MMT in air. Obviously, a full-scale separate study (well beyond the scope of this work) would be very important and should be undertaken.

The relative low volatility of MMT (i.e. its tendency to condense out on

surfaces) may also be a reason for the very low levels detected in air samples even in an underground car-park. With this in mind, the filter papers used for filtering particulate samples from the air in the MMT gas sampling train were rinsed and in some cases refluxed with isooctane. The rinsings were then injected into the gas chromatograph as a test for the presence of MMT, but in no case was MMT detected.

It is important to point out that tests done on exhaust samples taken from cars which were burning "unleaded" gasoline showed no MMT (detection limit 0.1 ng m^{-3}). This is consistent with a similar study of tetraalkyllead compounds [3]. Thus most of the MMT detected in the air samples is due to evaporation and spillage, etc.

This work was supported by a grant from the Ontario Ministry of the Environment, Research Grants Program and by Perkin-Elmer Corporation, Norwalk, Connecticut who supplied all the atomic absorption equipment. Thanks are also due to the Canada Center for Inland Waters, Burlington, Ontario, who lent the gas chromatograph.

REFERENCES

- 1 G. N. Turkel'traub, B. M. Luskina and S. V. Syavtsillo, *Khim. Tekhnol. Topl. Masel*, 12 (1967) 58.
- 2 P. C. Uden, R. M. Barnes and F. P. DiSanzo, *Anal. Chem.*, 50 (1978) 852.
- 3 B. Radziuk, Y. Thomassen, Y. K. Chau and J. C. van Loon, *Anal. Chim. Acta*, 105 (1979) 255.

THE INFLUENCE OF Mn_3O_4 FROM MMT COMBUSTION ON
GASOLINE VEHICLE EMISSIONS

Roy M. Harrison
Director
Institute of Aerosol Science
University of Essex
Colchester CO4 3SQ
United Kingdom

INTRODUCTION

In a trial of the influence of MMT fuel additive upon gasoline engine exhaust emissions, the Ethyl Corporation ran two matched fleets of vehicles, identically tuned and equipped with three-way catalyst systems. One fleet ran on fuel containing MMT additive at a concentration of 0.03125 g Mn per gallon of fuel, the other on an identical fuel without additive (termed "clear" fuel). The vehicles using MMT-additive showed statistically significant lower emissions of NO_x and CO, and slightly higher HC, than those running on clear fuel. The influence of MMT in reducing NO_x and CO emissions was greater over 0-75,000 miles than over 0-50,000 miles, whilst the effect on hydrocarbon emissions was marginally less over the greater distance.

This report seeks to explain the reasons for the observed differences in terms of the known catalytic properties of manganese oxides derived from MMT combustion.

SCIENTIFIC BACKGROUND

The combustion of MMT fuel additive in the gasoline engine leads to formation principally of the manganese oxide Mn_3O_4 ; traces of Mn_2O_3 have also been reported (1,2). Manganese oxides are well known for their catalytic properties, particularly in relation to their ability to oxidize carbon monoxide to carbon dioxide (3,4).

Past work in the laboratory (5) has shown that addition of MMT to iso-octane fuel in a pulse flame apparatus leads to reduced emissions of NO and CO, with increased hydrocarbons. The decrease in NO and CO emissions appeared to be associated with wall deposits of manganese oxides.

At the time of that work (1977/8), I formulated a series of experiments in conjunction with Professor Harry W. Edwards (Colorado State University). In these laboratory experiments, carried out at University of Lancaster (U.K.), the catalytic activity of Mn_3O_4 with respect to NO chemistry was investigated (6). We found that even in the presence of atmospheric oxygen, Mn_3O_4 catalyzed the decomposition of NO at temperatures as low as 215-243 degrees C (419-469 F). Concentrations of NO_2 were unaffected.

-2-

Nitric oxide, NO, is formed in the gasoline engine primarily by combination of atmospheric nitrogen, N_2 and oxygen, O_2 in a complex chain mechanism. The reaction is favored by high temperatures for two reasons. Firstly, the formation reaction is more rapid at elevated temperatures, and secondly the equilibrium concentration of NO is highly temperature sensitive with highest concentrations formed at high temperatures. The equilibrium constant, K, for the reaction is given by equation (1), from ref. (7):



$$K = \frac{X_{NO}}{X_{N_2}^{1/2} X_{O_2}^{1/2}} = 4.69 \exp (-21,600/RT) \quad (1)$$

Where X represents the partial pressure. At temperatures of 1000 K (1341 F) and below, the equilibrium concentrations of NO are very low. However, much higher concentrations can exist as the decomposition reaction is very slow at these temperatures and the NO is essentially "frozen in". A catalyst which speeds the decomposition reaction will lead to low-temperature breakdown of NO. It appears that Mn_3O_4 is such a catalyst (6). If 482 C (900 F) is taken as a typical exhaust gas temperature, the equilibrium partial pressure of NO may be calculated from equation (1). At an oxygen concentration of 1%, this is less than 1 ppm. Thus, in the presence of a suitable catalyst, NO decomposition would be expected.

INFLUENCE OF Mn_3O_4 IN THE VEHICLE EXHAUST

Scenario for NO Decomposition in Vehicle Exhaust

As a simplification, I have assumed that in a vehicle travelling at 30 mph, fuel of composition C_8H_{14} is combusted at a rate of 21 miles per U.S. gallon. MMT additive is used at a concentration of 0.03125 g Mn per U.S. gallon. The exhaust gases pass through a 2 meter length of 3 inch internal diameter pipe, at a mean temperature of 482 C (900 F). As it is known that only a small proportion of Mn burned in the engine is emitted in the exhaust, it is assumed that the walls of this pipe become coated with Mn_3O_4 .

Calculation of Influence Upon NO Concentrations

In my paper on NO decomposition over Mn_3O_4 , the reaction was found to be first order in NO with Arrhenius parameters of $E_{app} = 12.4 \pm 0.2 \text{ kcal mol}^{-1}$ and $\ln A(s) = 9.29 \pm 0.24$. Using the relationship,

$$k = A \exp (-E_{app}/RT) \quad (2)$$

-3-

(where R is the gas constant and T the absolute temperature), the rate coefficient k at 482 C (755 K) may be calculated to have a value of 2.93 s^{-1} .

In the plug flow reactor used in our study, the integrated rate expression was as follows:

$$-k V/u = \ln (C_e/C_i) \quad (3)$$

in which V = reactor volume
 u = volume flow rate
 C_e = steady state exit concentration
 C_i = steady state inlet concentration

In the vehicle exhaust scenario, the ratio V/u , the residence time, is calculated to be 0.24s.

Then, calculation gives

$$\frac{C_e}{C_i} = 0.50 \quad \text{at } 900 \text{ F}$$

This result implies 50% decomposition of NO and relates to conditions in our reactor. The precise surface area of Mn_3O_4 was not determined in our studies, but a minimum value was estimated as $0.005 \text{ m}^2 \text{ g}^{-1}$. As the charge was 4.03 g of Mn_3O_4 , the total surface area (minimum estimate) is 202 cm^2 . In the vehicle exhaust scenario, the area of exhaust pipe surface is $4,800 \text{ cm}^2$. The actual area of Mn_3O_4 deposits coating this surface will be appreciably greater as the surface deposit will have an uneven microstructure, and may be 10 fold, or more times this value. It is clear, therefore, that even if the specific surface area of Mn_3O_4 in our experiments is appreciably greater than the reported minimum value, the exposed surface area in the auto exhaust system is of a similar magnitude, assuming an even coverage of manganese oxide. Further decomposition of NO may also occur as a result of contact with Mn_3O_4 deposits in the three-way catalyst.

It is also possible to calculate the surface area of Mn_3O_4 aerosol in the exhaust gases. If spherical particles of diameter 10nm (typical of primary exhaust aerosol) are assumed, the surface area of aerosol present at any instant in the 2 meter length of exhaust is ca. 5 cm^2 , and thus much lower than that of the coated surfaces.

Surface-catalyzed reactions, at low conversions, typically show a mass conversion rate linearly proportional to the active surface area of the catalyst (3). At higher conversions in a plug flow reactor, a less than linear dependence would be expected, due to a reduction in

-4-

NO partial pressure as the gases pass through the reactor. The results of the calculation therefore lead me to the view that a coating of Mn_3O_4 in the vehicle exhaust could lead to a substantial reduction of the NO concentration discharged from the engine cylinder. This process will be accompanied by formation of O_2 .



Reductions of NO of the magnitude suggested above are quantitatively consistent with the observations of Otto and Sulak (5), who found a reduction in NO of 91% after 800 hours in experiments with the pulse flame apparatus with a constant level of CO.


Vehicle trials carried out by Ethyl Corporation have shown an increasing difference in emissions of both NO and CO with mileage between vehicles using clear fuel and fuel with MMT. This trend of improved performance in the MMT-fuel vehicles is consistent with the build-up of a surface coverage of Mn_3O_4 in the exhaust system.

Influence on Concentrations of CO

Our experiments did not include the influences of CO and HC. There is, however, substantial literature demonstrating the catalytic influence of manganese oxides, including Mn_3O_4 on oxidation of CO (e.g. ref.: 8)



Indeed, it is suggested (9) that Mn_3O_4 is the phase most active in this process. There are also precedents for the catalysis of CO oxidation by NO on metal oxides, including those of manganese (e.g. ref.: 10).



Whilst it is not possible to predict at this stage by which mechanism the effect occurs, it seems very probable that Mn_3O_4 is catalyzing CO oxidation and effecting an improvement in exhaust composition as a result. This effect was seen also in the work of Otto and Sulak (5) in the pulse flame apparatus. There are, to my knowledge, no detailed kinetic studies of these reactions available in the literature from which quantitative predictions might be made.

-5-

CONCLUSIONS

Operation of vehicles on fuel containing MMT additive causes deposition of a surface coating of the manganese oxide, Mn_3O_4 on the internal surface of the exhaust system. This coating has catalytic properties which cause decomposition of nitric oxide to N_2 and O_2 , and may also catalyze loss of carbon monoxide by reaction with O_2 , or with NO . This catalytic effect leads to an improvement in exhaust gas quality relative to vehicles running on clear fuel which increases with the age of the vehicle as the catalytic coverage of Mn_3O_4 grows.

REFERENCES

1. J. E. Faggan, A. D. Bailie, E. A. Desmond and D. L. Lenane, SAE Automobile Engineering Meeting Paper 750925 (1975)
2. EPA Report - 650/2-75-014 (1974)
3. P. G. Ashmore, Catalysis and Inhibition of Chemical Reactions, Butterworths (1963)
4. G. V. Samsonov, Oxide Handbook, Plenum, New York (1973)
5. K. Otto and R. J. Sulak, Environ. Sci. Technol., 12, 181-184 (1978)
6. H. W. Edwards and R. M. Harrison, Environ. Sci. Technol., 13, 673-676 (1979)
7. J. H. Seinfeld, Atmospheric Chemistry and Physics of Air Pollution, Wiley (1986)
8. J. W. Shulze et al., Z. Anorg. Allg. Chem. 452 67-76 (1979)
9. P. G. Tsyrlunikov et al., Kinet. Katal., 22, 1219-26 (1981)
10. T. G. Alkhazov et al., Kinet. Katal., 16, 1230 (1975)

013BFF90

APPENDIX

10



AN
ASSESSMENT OF THE EFFECT
OF MMT ON LIGHT-DUTY
VEHICLE EXHAUST EMISSIONS
IN
THE CANADIAN ENVIRONMENT

Prepared by a Working Group
Under the Jurisdiction of
The CGSB Gasoline and
Alternative Automotive
Fuels Committee

April 4, 1986

C O N T E N T S

SUMMARY

INTRODUCTION

OBJECTIVES

METHOD

Reports

Test Programs

Vehicle Manufacturers

MMT EFFECTS ON VEHICLE EMISSIONS

IMPACT ON FUTURE EMISSION CONTROL SYSTEMS

IMPACT ON AIR QUALITY

DISCUSSION

CONCLUSION

RECOMMENDATIONS

REFERENCES

BIBLIOGRAPHY

APPENDIX:

I Working Group

II Other Groups

III MVMA

IV AIC

V Ethyl Waivers To EPA

VI Volvo Data

SUMMARY

A working group of the CGSB Gasoline and Alternative Automotive Fuels Committee has reviewed the literature on the effect of manganese antiknock (MMT) on tailpipe emissions from light duty vehicles. It has concluded that the presence of MMT in commercial gasolines has no detectable effect on carbon monoxide or oxides of nitrogen emissions. Hydrocarbon emissions should increase on average from 0.03 to 0.11 g/mile due to the presence of MMT.

The emission-control systems developed in the U.S. and elsewhere to meet the present U.S. limit of 0.41 g/mile appear unlikely to suffer damage from exposure to MMT treated fuels. These emission-control systems will almost certainly be utilized on Canadian vehicles when this standard is adopted in Canada in 1988. A deterioration in HC control may be detectable over an 80,000 km (50,000 mile) test, but this would have no significant effect on overall average air quality. The major improvement (79.5%) will be due to the tightening of HC limits from 2.0 g/mile to 0.41 g/mile. Assuming the 0.11 g/mile increase, the continued presence of MMT in commercial gasolines would still permit a 74.0% improvement over the present standard.

On this basis, the working group recommends the retention of MMT at current levels in Canadian unleaded gasoline. Should changes in emission-control systems be introduced in the future, the question of their response to MMT should again be addressed.

★ ★ ★ ★ ★

INTRODUCTION

On August 3, 1985, the Department of Transport published, in the Canada Gazette Part 1, a notice proposing new emission standards for light-duty vehicles and light-duty trucks. The proposal would reduce the present allowable level of unburned hydrocarbon (HC) tailpipe emissions from 2.0 g/mile to 0.41 g/mile for light-duty vehicles and 0.8 g/mile for light-duty trucks, to take effect September 1, 1987, with the introduction of the 1988 model year. Vehicles will be tested for compliance with these standards using EPA procedures with accumulated mileage utilizing MMT-free gasoline.

In adopting these tighter standards, the use of the octane enhancer methylcyclopentadienyl manganese tricarbonyl (MMT*) in commercial pump gasoline would be continued at the current Canadian General Standards Board (CGSB) limit of 18 mg Mn/litre unless its continued use could be shown to have unacceptable adverse effects on health or on the environment.

There are sound economic reasons for the retention of MMT in Canadian unleaded gasolines. This useful antiknock agent saves energy and money in meeting the octane requirements of the Canadian automobile population. However, the effects of MMT on the Canadian environment, with particular reference to the new standard, have not been fully evaluated.

To assess the potential effects of manganese compounds on vehicle emissions and on emission system durability in Canada, Environment Canada requested assistance from the Canadian General Standards Board Petroleum Committee. The CGSB Gasoline and Alternative Fuels Committee subsequently appointed a working group to carry out this activity. The working group comprised members from the petroleum industry, the vehicle manufacturers and the federal government (Appendix I). Comments and advice were solicited from other interested parties such as health and consumer groups (Appendix II).

OBJECTIVES

The objectives of the working group, outlined in a letter (dated 27/5/85) by the Assistant Deputy Minister, Environmental Protection Service, Dr. R.W. Slater, to Dr. R. Whyte, Chairman of the Petroleum Committee, were as follows:

- i) to review and report on the current state of knowledge respecting MMT effects on vehicle emissions,

*Ethyl Corporation's TM

- ii) to forecast the possible impact on the vehicle emission control systems likely to be in use in the period 1988-2000,
- iii) to assist Environment Canada in documenting the impact that continued use of MMT may have on air quality.

This report presents the findings and judgments of the working group.

METHOD

To accomplish this task, the working group gathered the information through the following methods.

(i) Reports

Relevant reports on MMT were compiled and reviewed (see bibliography). It was noted that very little work had been reported since 1978 when the use of MMT in unleaded gasoline was disallowed by the U.S. Environmental Protection Agency (EPA).

(ii) Test Programs

The results of two test programs were reviewed by the working group; the "CRC MMT Field Test Program" reported in June 1979; and a recent in-use vehicle surveillance test program conducted by Environment Canada which tested 1983 to 1985 model year Cadillacs, calibrated to U.S. emission specifications (References 1 and 2).

(iii) Vehicle Manufacturers

Information was requested from both the Motor Vehicle Manufacturers Association (MVMA) and the Automobile Importers of Canada (AIC) member companies (Appendix III and IV), as to how current emission-control systems were behaving in Canada and how future emission-control systems would behave if MMT were retained in gasoline at the current CGSB limits.

MMT EFFECTS ON VEHICLE EMISSIONS

The working group realized from the outset that very few studies had been reported on MMT and its effects on exhaust emissions, since it was disallowed by the EPA in the U.S. for use in unleaded gasoline in 1978. Subsequent applications for a waiver by Ethyl Corporation to use MMT in U.S. unleaded gasolines were not granted (Appendix V).

An extensive test program was carried out by the Co-ordinating Research Council (CRC) in 1977-78, and more recently Environment Canada conducted an in-house study, both of which have been extensively reviewed by the working group.

1) CRC MMT FIELD TEST PROGRAM

The CRC program involved 63 vehicles. The objective of the CRC program was to study the effects of two concentrations of manganese on exhaust emissions and, to compare these results with those for clear fuel. The 63 vehicles were divided into three matched fleets. The first fleet was operated on clear gasoline, the second on gasoline which contained MMT at a level of 1/32 gMn/USG (8.25 mgMn/L), and the third, on gasoline which contained MMT at 1/16 gMn/USG (16.5 mgMn/L). Within each fleet, seven car models were tested in triplicate. In other words, nine replicate cars of each of the seven car models were tested resulting in a 63-car test program.

Of the seven car models studied in the CRC program, only two featured three-way catalysts (TWC) and closed-loop fuel systems, the types likely to be in use in Canada by the 1988 model year. These two models were a Pontiac Sunbird and a Volvo 242, the former being electronic feedback carbureted, and the latter having feedback electronic fuel injection. Five car models featured conventional oxidation catalysts (COC) and open-loop fuel systems. All of these vehicles were essentially production models which had been calibrated to meet 1977-78 California emission standards of 0.41 g/mile HC, 9.0 g/mile carbon monoxide (CO) and 1.5 g/mile oxides of nitrogen (NO_x). These vehicles had been selected by their respective manufacturer since they were considered to be sensitive to MMT and/or they were representative of future high-volume powertrains.

The presence of MMT was found on average to be statistically significant only for HC, and did not affect CO or NOx emissions. While the vehicles exhibited varying initial HC emissions, all were calibrated to meet a 0.41 g/mile hydrocarbon standard.

The study found that, on average for all vehicles, the tailpipe HC emissions increased by 0.11 g/mile relative to clear fuel, after the 50,000-mile test period while using MMT at 1/16 gMn/USG (16.5 mg Mn/litre) in the fuel.

The data from the two TWC model cars were not consistent, and ranged from zero to 0.17 g/mile difference in HC emissions relative to clear gasoline at 50,000 miles. The Volvo cars, produced the lowest HC emissions and had the smallest response to MMT. The average HC increase relative to clear gasoline was 0.03 g/mile when using 1/16 g Mn/USG (Appendix VI - Composite Results Prepared by the Working Group). The applicability of this result to all vehicles equipped with fuel injection and TWC is questionable since TWC electronic fuel-injection control system may not be more efficient than a COC system at controlling the increased HC emissions due to MMT which are generated in the combustion chamber. Electronic fuel injection will allow better control of the air-fuel ratio, and this will only reduce the 'spread' of HC emissions when compared to carbureted vehicles. However, of the cars tested in the study, the Volvo fuel and emission-control system was closest to that likely to be sold in Canada during the 1988 model year and beyond.

In addition, there was no indication of catalytic converter plugging with any of the fuels in the test program, as monitored by the pressure drop across the catalyst. In no instances was the fuel economy of the vehicles of the CRC test fleet affected by the MMT in the gasoline.

The CRC study also found that the use of MMT resulted in a catalytic converter efficiency increase of 2 to 3%, although this was not enough to offset increases in engine-out emissions.

2) ENVIRONMENT CANADA STUDY

In 1984 Environment Canada, in order to generate HC emission data, conducted surveillance tests of in-use vehicles calibrated to U.S. emission control specifications which had been operating on Canadian unleaded gasoline containing MMT at levels of up to 18 mg Mn/L. The cars, model years 1983, '84 and '85, featured electronic throttle-body fuel-injection, an oxidation catalyst working in conjunction with a three-way catalyst, and closed-loop fuel control. Tests were not undertaken on misfueled vehicles or vehicles with malfunctioning fuel-control systems. Fifteen rented in-use Cadillac cars equipped with 4.1-litre engines were tested and sixteen acceptable Federal Test Procedure (FTP) tests were performed.

The results indicated that, when using data generated by a different vehicle at each test point, these cars would not exceed the proposed hydrocarbon emission level of 0.41 g/mile when test results were extrapolated to 50,000 miles. However, any conclusion regarding HC emission increase, if any, due to MMT in the gasoline, could not be established due to the absence of base-line data for similar vehicles operated on clear fuels.

The major conclusions based on these two test programs may be summarized as follows:

1. Based on all cars tested in the CRC study, the average increase in tailpipe HC emissions at 50,000 miles using MMT at 1/16 g Mn/USG was 0.11 g/mile higher than for clear fuel. This may be taken as the best average case.
2. Based only on the two TWC car tests in the CRC study, the average HC increase for the 1/16 g Mn/USG fuel ranged from zero to 0.17 g/mile higher than for clear fuel, at 50,000 miles.
3. For the Volvo system, which was assumed to be closest to the current and future technology, the average HC increase relative to clear fuel was 0.03 g/mile (Working group analysis of Volvo data). This may be taken as the best case.

4. All cars in the Environment Canada study met the proposed HC standard of 0.41 g/mile on fuel containing MMT, when extrapolated to 50,000 miles. The effect of MMT on vehicle HC emissions was inconclusive since no clear fuel base line data were obtained. However, there were no indications that the MMT effect is greater than that indicated in the CRC report.

5. The effect of MMT on HC emissions may be of the same order of magnitude as the test uncertainty in a small fleet test. Therefore, it may not be possible to measure statistically significant differences due to MMT, especially in cases where the majority of cars meet the HC standard.

IMPACT ON FUTURE EMISSION-CONTROL SYSTEMS

VEHICLE MANUFACTURERS

The MVMA and AIC have been asked to predict how future emission-control systems would behave if MMT were retained in unleaded gasoline. Based upon limited data available, there is no evidence to suggest that emission-control systems that have been developed for today's market would experience adverse effects with regard to function or performance from the use of MMT in gasoline. Current systems do not indicate that durability is lower in Canada, where MMT is used, than in the United States where MMT has been disallowed in unleaded gasoline. Members of MVMA and AIC indicate that manufacturers' Canadian warranty claims on emission components are comparable to the U.S.

It must be noted however that the effect of MMT on emission-control system components may be subtle and may never result in a failure which would be apparent through analysis of warranty claims. Degradation of the emission performance of a vehicle does not necessarily manifest itself in the form of poor driveability or performance yet; when emission levels above those may display an increase in emission levels above those to which the vehicle was designed.

The durability of critical emission-control components related to MMT use may change in the future. This may be reflected in the manufacturers' warranty records. Therefore, short of complete emission measurements, periodic reviews of warranty records may be necessary to determine if future action on MMT is required.

In the past, some researchers noted that MMT tended to form deposits, typically manganese oxide, which coated the oxygen sensor in the exhaust system of closed-loop type cars. Depending upon the design of the oxygen sensor, deposition of manganese oxide could impede the sensor's ability to function properly in controlling exhaust emissions, as was evidenced in the CRC program already cited. A failure of this nature may or may not be accompanied by a gross deterioration in vehicle driveability.


The working group has considered that further test programs on vehicles using today's equipment could well provide additional information. However, this working group has received no evidence to suggest that an average increase would be greater than 0.11 g/mile. Future technologies suggest that exhaust emission-control systems will be more effective at controlling emissions sensitive to the effects of MMT, and therefore further large-scale testing may be of limited value.

The working group would recommend a test program only in the event that future emission-control technology was demonstrated to be adversely affected by MMT. The CGSB Committee on Gasoline and Alternative Automotive Fuels would be receptive to any new data generated on MMT fuels.

GENERAL DISCUSSION

The effect of MMT usage on HC emissions is shown in the graph below. The HC emission reductions since 1971 have also been indicated.

Canadian Hydrocarbon Emission Standards



COMMENTS

- 1) The proposed standard of 0.41 g/mile represents a reduction of 79.5% from the present standard of 2.0 g/mile.
- 2) Assuming that MMT increases HC emissions on average by 0.11 g/mile relative to clear fuel, an actual HC level of 0.52 g/mile ($0.41 + 0.11$) represents an average reduction of 74.0% from the present standard.
- 3) Assuming that MMT increases HC emissions by 0.03 g/mile relative to clear fuel, an actual HC level of 0.44 g/mile ($0.41 + 0.03$) represents an average reduction of 78.0% from the present standard.

The actual average increase in HC tailpipe emissions should lie somewhere between these two values and consequently the working group cannot see any significant benefit in removing MMT from unleaded gasoline in Canada based on tailpipe HC emission levels.

IMPACT ON AIR QUALITY

The working group has not attempted to carry out any very rigorous study on the effect of retaining MMT on overall air quality. Environment Canada have, however, published a "National Inventory of Natural Sources and Emissions of Organic Compounds" in which the total man-made emissions of hydrocarbons are estimated as 8.4% of the whole (Reference 3 and 4). Of these, 16% are attributed to light-duty vehicles.

It can thus be calculated that reducing the hydrocarbon standard from 2.0 g/mile to 0.41 g/mile (79.5%) will result in a reduction in HC burden of $79.5 \times 0.16 \times 0.084 = 1.07\%$.

In the worst case, an increase of 0.11 g/mile due to MMT, the overall reduction would be 74%. This equates to an improvement in air quality of 0.99%. In the best case, an increase of 0.03 g/mile due to MMT, the emission reduction would be 78%, equivalent to an improvement in air quality of 1.05%.

NATIONAL ENVIRONMENTAL INVENTORY SYSTEM (NEIS)

The level of HC tailpipe emission increase due to MMT will be required by the Department of the Environment in order to calculate an emission factor for use in NEIS. Analysis of the data has indicated that in future, with the majority of new cars being fitted with three-way catalysts, fuel injection and closed-loop fuel control systems, the average increase in tailpipe HC emissions could be as high as 0.11 g/mile for post-1988 model cars. It is recommended that this value be used by the Department of the Environment to calculate the emission factor.

CONCLUSIONS

The major conclusions of the working group based on the analysis of currently available data are as follows:

- 1) When using MMT the average increase in tailpipe HC emissions over clear fuel should range from 0.03 to 0.11 g/mile for the 1988 model year light-duty vehicles.
- 2) The proposed standard would reduce the allowable tailpipe HC emissions from 2.0 g/mile to 0.41 g/mile. With clear fuel this is a HC reduction of 79.5%. With MMT the average reduction would be from 74% to 78%.
- 3) The use of MMT at current CGSB levels does not significantly compromise emission-control system operation or component durability.
- 4) The working group has not attempted to carry out any rigorous study on the effect of retaining MMT on overall air quality. However, since it is estimated that vehicle emissions contribute only 1.35% of hydrocarbons present in the atmosphere, the effects of MMT is considered miniscule.

RECOMMENDATIONS

- 1) Based on the findings in this report, the members of the working group recommend that MMT be retained at current levels as an octane enhancer in unleaded gasoline.
- 2) The members of the working group recommend that the issue be re-examined in the event that future emission-control technology or the generation of additional data, should show adverse effect due to the use of MMT beyond those recognized in this report.

Signed.....
Chairman of Working Group

REFERENCES

- 1) COORDINATING RESEARCH COUNCIL, INC.,
"CRC MMT Field Test Program", 503, June 1979.
- 2) OSTROUCHOV, N.;
"MMT Effect on Hydrocarbon Emissions"
Environment Canada, Memorandum, Ottawa, Ontario,
March 21, 1985.
- 3) ENVIRONMENT CANADA:
"National Inventory of Natural Sources and
Emissions of Organic Compounds."
- 4) PACE
Comments on Proposed 1986 Light Duty
Automotive Emission Standards.

BIBLIOGRAPHY (1 of 4)

MMT BIBLIOGRAPHYPAPERS

ANON; "MMT Revisited: EPA's conclusions"; Automobile Engineering Volume 87 Number 6; June 1979.

BAILIE, J.D.; "Economics of Manganese as an Antiknock in Unleaded Gasoline"; NPRA Annual Meeting, March 1976; AM-76-17.

BAILIE, J.D.; "MMT: A Better Way to Boost Octane"; Ethyl Corporation, Houston Texas, Oil Gas Journal, Volume 74, Number 16; April 19, 1976.

BAILIE, J.D.; "The Status of MMT"; NPRA F&L meeting; November 1977, F&L-77-66.

BAILIE, J.D., Desmond, E.A.; Fagan, J.E.; Leman, D.L.; "An Evaluation of Manganese as an Antiknock in Unleaded Gasoline"; SAE 750925, Automobile Engineering Meeting, Detroit, Michigan, October 13, 1975.

BAILIE, J.D.; Michalski, G.W.; Unzeman, AG.H.; "MMT Outlook 1977"; Ethyl Corporation, Baton Rouge, LA.; API (Volume 56; Order Number 821-01977), Washington, DC.; 1977.

BENSON, J.D.; Campion, R.J.; Painter, C.J.; "Results of Coordinating Research Council MMT Field Test Program"; SAE 790706, Passenger Car Meeting, Hyatt Regency, Dearborn; June 11-15, 1975.

BOORAS, S.G.; "Airborne Manganese Health Effects and Recommended Standard"; Environmental Health Research Center; July 1975.

BRADCOCK, J.N.; Duncan, J.; "Combustor Study of the Deactivation of a Three Way Catalyst by Lead and Manganese"; SAE 841408, Fuels and Lubricants Meeting and Exposition, Baltimore; October 8-11, 1984.

BUMBACO, M.; "Methanol Development - MMT in Gasoline"; EPS-Analytical Services Division, Technical Services Branch, Ottawa, Ontario, May 11, 1984.

CALABRESE, E.J.; Sorenson, A.; "Methylcyclopentadienyl Manganese Tricarbonyl as an Antiknock Component: Composition and Fate of Manganese Exhaust Products", University of Illinois at the Medical Center, Chicago; APCA Journal, Volume 25, Number 12, December 1975.

CARLSON, R.R.; Pilkington, R.E.; Politzer, J.L.; "Conduct of Mileage Accumulation and Emission Testing for the CRC/MMT Program"; SAE 790703, Passenger Car Meeting, Hyatt Regency, Dearborn; June 11-15, 1979.

Bibliography (2 of 4)

COOPER, L.W.; "The Health Implications of Increased Manganese in the Environment Resulting from the Combustion of Fuel Additives: A Review of the Literature"; Journal of Toxicology and Environmental Health, 14:23-46, 1984.

COORDINATING Research Council, Inc., "Study of the Effects of MMT on Automotive Exhaust Emissions", EPA Hearing on MMT Waiver Request, Washington, D.C., June 27-30, 1978.

DONALDSON, J.; LaBella, F.S.; Gresser, D.; "Enhanced Auto-oxidation of Dopamine as a Possible Basis of Manganese Neurotoxicity"; University of Manitoba, Winnipeg, Manitoba, 1980.

"EPA Test Plan to Study the Effect of MMT on Emission Control System Performance - Plan Summary", (undated).

FAIX, L.J.; "A Study of the Effects of Manganese Fuel Additive on Automotive Emissions"; SAE 780002, Congress and Exposition, Cobo Hall, Detroit; February 27 - March 3, 1978.

FERN, G.R.H.; "PACE Estimated Refinery Investment for New LDV Emission Standards"; Petroleum Association for Conservation of the Canadian Environment; Ottawa, Ontario; July 20, 1984.

FUREY, Robert L.; Summers, Jack C.; "How MMT Causes Plugging of Monolith Converters", SAE 780004, Congress and Exposition, Cobo Hall, Detroit, February 27 - March 3, 1978.

GANDHI, H.S.; Weaver, E.E.; Williamson, W.B.; "Effects of Fuel Additive MMT on Contaminant Retention and Catalyst Performance"; SAE 821193, Fuels and Lubricants Meeting, Toronto, Ontario; October 18-21, 1978.

GARBE, Robert J.; "Minutes of January 20, 1977 MMT Meeting", U.S. EPA, Motor Vehicle Emission Laboratory, Ann Arbor, February 14, 1977.

GARBE, Robert J.; "Minutes of March 29, 1977 MMT Meeting", U.S. EPA, Briarwood Hilton, Ann Arbor, April 20, 1977.

GARBE, Robert J.; Wallace, Jay S.; "Effects of MMT on Exhaust Emissions", SAE 790707, Passenger Car Meeting, Hyatt Regency, Dearborn, June 11-15, 1979.

HEALTH and Welfare; "MMT - An assessment of the Human Health Implications of its Use as a Gasoline Additive", Report No. 78-EHD-21, 1978.

HIGGINS, P.M.; "MMT (methycyclopentadienyl manganese tricarbonyl) and HC (hydrocarbon) Emissions from Light Duty Motor Vehicles"; EPS - Transportation Systems Division Memorandum, Ottawa, Ontario.

Bibliography (3 of 4)

HOLMES, J.R.; "Health Effect Evaluations of Manganese and Particulate Matter"; EPS September 21, 1984.

HUGHMARK, C.A.; Sobel, B.A.; "A Statistical Analysis of the Effect of the MMT Concentration on Hydrocarbon Emissions"; SAE 800393, Congress and Exposition, Cobo Hall, Detroit, February 25-29, 1980.

INDUSTRIAL Incentives Division, Technical Services Branch, "MMT in Lead Free Gasoline"; EPS Ottawa, Ontario, September 13, 1984.

JOST, K.C.; Gordon, C.E.; "Safe Handling of MMT in Refineries"; Ethyl Corporation, Houston, Texas, API (Volume 56, Order Number 821-01977); 1977.

LANDHEER, F.; "Monenco - MMT Study"; Transportation Systems Division Memorandum, September 5, 1984.

LENANE, D.L.; "MMT - A Further Evaluation", SAE 770656; F&L Meeting, June 1977; Tulsa, OK.

LENANE, D.L.; "Effect of MMT on Emissions from Production Cars", SAE 780003, Congress and Exposition, Cobo Hall, Detroit, February 27 - March 3, 1978.

LICHTENSTEIN, Ivan E.; Mundy, Joseph P.; "MMT Plugging of Oxidation Catalysts on Ceramic and Metal Supports During Engine Dyno Studies of Catalyst Durability", SAE 780005, Congress and Exposition, Cobo Hall, Detroit, February 27 - March 3, 1978.

MOORE, W.; Stara, J.F.; Hysell, D.; Malanchuck, M.; Burkhart, J.; Hinnners, R.; "Toxicologic Evaluations of Fuel Additive - Methylcyclopentadienyl Manganese Tricarbonyl (MMT)", SAE 750927, Automobile Engineering Meeting, Detroit, Michigan, October 13-17, 1975.

MORAN, J.B.; "The Environmental Implications of Manganese as an Alternate Antiknock"; SAE 750926, Automobile Engineering Meeting, Detroit, Michigan, October 13-17, 1975.

OSTROUCHOV, N.; "MMT Effect", EPS - Transportation Systems Division Memorandum, Ottawa, Ontario, November 16, 1984.

OSTROUCHOV, N.; "MMT Effect on Hydrocarbon Emissions" EPS - Transportation Systems Division Memorandum, Ottawa, Ontario, March 21, 1985.

OTTO, K.; Sulak, R.J.; "Effects of Mn Deposits from MMT on Automotive Catalysts in the Absence and Presence of Other Fuel Additives", Ford Motor Company, American Chemical Society, Volume 12, Number 2; February 1978.

Bibliography (4 of 4)

PIERSON, W.R.; McKee, D.E.; Brachaczek, W.W.; Butler, J.W.;
"Methylcyclopentadienyl Manganese Tricarbonyl: Effect on Manganese Emissions
from Vehicles on the Road"; Ford Motor Company, APCA Journal 1978.

STATE of California Air Resources Board, "Manganese Fuel Additive MMT", Staff
Report 77-9-3, April 28, 1977.

WALLACE, J.S.; Garbe, R.J., "Effect of MMT on Exhaust Emissions"; SAE 790707,
Passenger Car Meeting, Hyatt Regency Dearborn, June 11-15, 1979.

WOTZAK, Gregory P.; "Catalyst Plugging in Thruway Police Vehicles - Interim
Report", New York State Department of Environmental Conservation Presentation
to EPA Conference on MMT Projects, Ann Arbor, March 29, 1977.

UNZELMAN, G.H.; "Antiknock for Unleaded Gasoline - Economic, Energy and
Environmental Aspects", Western Gas Processors and Oil Refiners Association,
October 1975.

U.S. Federal Register; "Decision Not to Regulate Manganese Under the Clean Air
Act:", August 13, 1985 p 32627.

* * * * *

LETTER

LEWIS, William H., State of California Air Resources Board, "California Air
Resources Board Resolution Regarding Use of MMT", May 6, 1977.

* * * * *

TECHNICAL REPORTS

ETHYL Corporation Research Laboratories; "Effect of 'Ethyl' MMT on Soot
Fouling of Spark Plugs", ERM 76-1, February, 1976.

ETHYL Corporation, Ferndale, Michigan; "MMT Waiver Request", Updated
Report 1 and 2, June 27, 1978.

* * * * *

APPENDIX ICGSB WORKING GROUP ON MMT IN GASOLINE

<u>NAME</u>	<u>AFFILIATION</u>
K. Miller	Petro-Canada Products Inc.
R. Solman	Environment Canada
P. Sarvos	Shell Canada Ltd.
G. Caldwell	Motor Vehicle Manufacturers' Association
R. Falkiner	Esso Petroleum Canada
J. Collins	Ethyl Canada Inc.
J. Dowell	Energy, Mines and Resources
M. Mokry	Automobile Importers of Canada

The Canadian Association
for Children and Adults
with Learning Disabilities



L'Association canadienne
pour enfants et adultes ayant
des troubles d'apprentissage

Maison Kildare House, 323 Chapel, Suite 101, 102, Ottawa, Ontario K1N 7Z2 (613) 238-5721

August 27, 1985

Mr. J.O. Cliffe, Chairman
C.G.S.B. Committee on Gasoline & AAF
432 Martin Grove Road
Islington, Ontario
M9B 4M2



Dear Mr. Cliffe:

Thank you for your letter and attached information concerning the working group under the Canadian General Standards Board investigating the long-term effects of the continued use of MMT in unleaded gasoline.

I'm sorry I was unable to contact you or attend, but I was out of the province in July on vacation. As you know the conclusion stated in the report IP-8 The Technology and Costs of Control Automobile Emissions in Canada is as follows: "The consensus of EPA and the automotive industry is that MMT produces significant adverse effects on HC emissions." The authors were especially concerned about failure of sensors and other effects on the catalyst.

Our primary concern in the long-term use of MMT is with health, particularly its neurotoxic attributes in the case of young and unborn children. I have communicated these concerns directly to Mr. Hazra of EPS, and Dr. Joseph Ruddick of the Health Protection Branch of Health and Welfare.

Sincerely yours,

Barbara McElgunn
Research and Liaison Officer (Health)

APPENDIX 11
(Page 2 of 3)

432 Martin Grove Road,
Islington, Ontario
M9B 4M2

September 3, 1985

Mrs. Barbara McElgunn,
The Canadian Association for
Children & Adults with
Learning Disabilities,
Kildare House,
323 Chapel, Suite 101, 102
Ottawa, Ontario
K1N 7Z2

Dear Mrs. McElgunn:

Thank you for your letter of August 27th concerning the Working Group of C.G.S.B. set up under the auspices of my committee to review the data available on MMT in gasoline and its effects on automobile emissions. I have passed it on to Mr. Keith Miller of PetroCanada who is the chairman of the Working Group.

I was, of course, aware of your earlier contact with Mr. Hazra since it was he who had given me your address.

I understand that the Working Group has held one meeting since the inaugural one and will be meeting again sometime in October. By copy of this letter, I am asking Mr. Miller to ensure that you remain up to date on any meeting notes which may have been issued. If you need to contact him, his phone number is 416-445-9113. I will, myself, be out of town most of October and the first two weeks of November.

Yours very truly,



J.O. Cliffe

cc. Mr. K. Miller,
PetroCanada Products Inc.,
240 Duncan Mill Road,
Don Mills, Ontario
M3B 3B2